

PUBLIKASI PADA JURNAL INTERNASIONAL BEREPUTASI

“JOURNAL OF TROPICAL LIFE SCIENCE”

**“Nutritional Content and Characteristics of Antimicrobial
Compounds from Fermented Oil Palm Fronds (*Elaeis guineensis*
Jacq.)**

Nama : Dr. Dewi Febrina. S.Pt.,

NIP : 19730202 200501 2 004

NUPTK : 2534751652231152

ARTIKEL YANG DISUBMIT

Nutritional Content and Characteristics of Antimicrobial Compounds from Fermented Oil Palm Fronds (*Elaeis guineensis* Jacq)

Dewi Febrina¹, Rahmi Febriyanti², Syukria Ikhsan Zam^{2*}, Zumarni¹, Jepri Juliantoni¹,
Abdul Fatah¹

¹ Department of Animal Science, Faculty of Agriculture and Animal Sciences, State Islamic
University of Sultan Syarif Kasim Riau, Pekanbaru 28293, Indonesia

² Department of Agrotechnology, Faculty of Agriculture and Animal Sciences, State Islamic
University of Sultan Syarif Kasim Riau, Pekanbaru 28293, Indonesia

* corresponding author : email : syukria.ikhsan.zam@uin-suska.ac.id

ABSTRACT

This research was conducted to determine the nutritional quality and characteristics of antimicrobial compounds from Fermented Oil Palm Fronds (FOPOF). This research was carried out using a Complete Randomized Design with 4 treatments and 4 replications. The following treatments were performed the addition of different additives namely : 10% poultry manure, 10% cow feces, 5% urea and 5% molasses. The data were analyzed using a one-way analysis of variance and differences between treatments were tested using Duncan's multiple range test. The observed parameters were physical quality tests include pH, aroma, color , texture and nutritional quality including the content of NDF, ADF, hemi cellulose, cellulose and lignin. Extraction of oil palm fronds using multilevel maceration method with ethyl acetate, n hexane and methanol. Identification is done through phytochemical screening. The activity test of antimicrobial compounds extract from FOPF using paper disc method with a concentration of 1-5. FOPF with poultry manure yields the lowest pH value, sour aroma, yellowish green, softer texture and the lowest lignin content (19.94%). Methanol extract from FOPF with poultry manure produces the highest extract of rendemen and the largest inhibition zone diameter with a MIC is 1%.

Keywords : oil palm frond, extract methanol, antimicrobial, E. coli, S. aureus

Background

The utilization of oil palm plantation waste such as oil palm fronds as feed is still limited due to the high lignin content [1-2]. Physical-chemical-biological and combination treatments and the addition of additives on the fermentation process aim to break down lignocellulose and ligno hemicellulose bonds and increase nutrient digestibility [3-5].

Palm oil fronds have antimicrobial activity [6]. Methanol extract from palm oil leaves contains tannin, alkaloids, saponins, terpenoids

and flavonoids [7]; and toxicity tests on shrimp and rats showed that palm oil leaf extract is safe for consumption and recommended as a natural product [8]. This study aims to increase the nutritional content of oil palm fronds through a fermentation process with the addition of different additives and to determine the bioactive compounds in extracts from FOPF with multilevel extraction using n hexane, ethyl acetate and methanol.

Methods.

The research was conducted in June-November 2017, at the Laboratory of Nutrition and Chemistry at the Faculty of Agriculture and Animal Sciences of State Islamic University of Sultan Syarif Kasim Riau Indonesia, Biotechnology Research Center of the Indonesian Sciences Institute Indonesia and Laboratory of Biota Sumatra Andalas University.

Materials

The materials used in this research are NDS solution, ADS solution, acetone, alcohol, octanol, H₂SO₄, methanol, ethyl acetate, hexane, filter paper, rotary vacuum evaporator, HCl, ammonia, mayer reagent, dragendorff reagent, ether, Liebermann Bouchard reagent, Mg powder, amyl alcohol, NaOH, FeCl₃, aquadest, poultry manure, cow feces, urea, molasses, plastic bags. The tools used in the study were leaf chopper, crusibel dish, fibertec hot extraction, fibertec cold extraction, oven, desicator, furnace, analytical scales, dropple plate, water bath.

Research Procedure

The study was conducted through two (2) stages : namely fermentation of oil palm frond

based on Febrina [9] modified with different additive (10% poultry manure, 10% cow feces, 5% urea and 5% molasses) using a Completely Randomized Design 4 treatment with 4 replications. Fermentation process for 21 days. The data were analyzed using a one-way analysis of variance and differences between treatments were tested using Duncan's Multiple Range Test. The observed parameters were physical quality tests include pH, aroma, color and texture and nutritional quality including the content of NDF, ADF, hemi cellulose, cellulose and lignin based on Van Soest analysis using the FOSS Fibertec system. The second stage is multilevel maceration and extraction using methanol, ethyl acetate and n hexane solvents. Phytochemical screening based on Franswort [10] and the antibacterial activity test of oil palm fronds extract based on Handayani et al [11].

Results and Discussion

The effect of different additives on the physical quality of FOPF

The addition of different additives has a significant effect (P<0.05) on the physical quality of FOPF including pH, aroma, color and texture (Table 1).

Table. 1 The effect of different additives on the physical quality of FOPF

Treatment	Physical quality							
	pH		Aroma		Color		Texture	
	Value	Notes	Value	Notes	Value	Notes	Value	Notes
A	5,18 ^a ±0,10	Acid	3,00 ^b ±0,00	Acid	3,03 ^a ±0,05	yellowish green	2,00 ^a ±0,00	Rather soft
B	5,52 ^b ±0,20	Acid	4,00 ^c ±0,00	more acidic	3,00 ^a ±0,00	yellowish green	3,00 ^b ±0,00	soft
C	7,60 ^c ±0,14	Base	2,03 ^a ±0,05	Non acid	4,03 ^b ±0,05	brownish green	3,03 ^b ±0,05	soft
D	5,28 ^{ab} ±0,21	acid	5,00 ^d ±0,00	Very acid	3,00 ^a ±0,00	yellowish green	3,00 ^b ±0,00	soft

Notes : A = FOPF with the addition of poultry manure
B = FOPF with the addition of cow feces
C = FOPF with the addition of urea
D = FOPF with the addition of molasses

a,b,c means with different superscript differ ($p < 0.05$)

The lowest of pH (5.18) in treatment A is related to the availability of carbohydrates and the production of Lactic Acid Bacteria. The same results were seen in treatment D with a pH of 5.28. Addition of poultry manure and molasses on fermentation oil palm frond aims as a source of additive which can trigger the growth of Lactic Acid Bacteria to produce lactic acid. Increased production of lactic acid will reduce pH and inhibit the growth of pathogenic bacteria. Ozduven et al [12] stated that pH is associated with lactic acid produced during the fermentation process which can inhibit the growth of pathogenic microbes. Oil palm fronds which are fermented with molasses produce pH 3.93 [3].

The process of ammonization of oil palm fronds using urea will produce the highest pH of 7.60 which is alkaline. The same results were reported by Zahari et al [3], the process of ammonization of oil palm fronds using urea resulting in a pH of 7.38. In the process of ammonization with urea will break down into ammonium ions so that the ammonization product is alkaline. Laura and Paolo [13] state that urea will break down into ammonium ions, which are alkaline and ammonization processes can break down glycoside bonds [14] so that the oil palm fronds can be degraded by rumen microbes.

In treatment A produces an aroma score of 3 with acid criteria. This shows that the fermentation process is good which is characterized by acidic products were typical of fermentation. Aroma is related to the production of lactic acid and pH. It can be seen in Table 1 the addition of poultry manure to oil palm fronds fermentation resulting in the lowest pH (5.18) so that the products are acidic.

Treatment A, B and D produces a color that is not different ($P > 0.05$), which is natural green or yellowish green with score of 3-3.03. Good quality silage will produce colors that are not different from the original material [15]. The occurrence of color changes from green to yellowish green due to the anaerobic phase oxygen has run out then carbohydrates will produce CO_2 , H_2O and heat. The heat produced causes a change in color from green to yellowish green.

Treatment A will produce softer textures with score 2. This shows that the addition of the poultry manure to the fermentation process can support the growth of Lactic Acid Bacteria to produce lactic acid. Optimal production of lactic acid will reduce pH so that it inhibits the growth of pathogenic bacteria. This low pH also supports the production of lignase enzymes which function to break down lignocellulose and lignohemicellulose thereby reducing lignin content. It is seen that treatment A produced the lowest lignin content of 19.94%. The release of lignin from cellulose and hemicellulose causes fermented products to have a softer texture.

The effect of different additive on fiber fraction of FOPF

The addition of different additive had no effect ($P > 0.05$) on the content of NDF, ADF, hemicellulose and cellulose but had a very significant effect ($P < 0.01$) on the lignin content. Rahman et al [4] reported there was a decrease in the lignin content of oil palm fronds in the fermentation process using white rot fungi.

Table 2. The effect of different additive on fiber fraction of FOPF.

Treatment	Fiber Fraction (%)				
	NDF	ADF	Hemicellulose	Cellulose	Lignin
A	73,34±3,68	51,81±4,82	21,52±6,07	28,54±3,36	19,94 ^a ±1,47
B	75,39±2,03	55,03±4,23	20,35±4,46	29,57±3,09	23,15 ^b ±0,96
C	73,83±1,01	53,30±3,43	20,51±4,30	31,55±1,99	21,57 ^a ±1,58
D	74,98±3,52	54,04±4,48	20,93±6,63	31,70±4,27	22,41 ^b ±1,22

Notes: A = FOPF with the addition of poultry manure

B = FOPF with the addition of cow feces

C = FOPF with the addition of urea

D = FOPF with the addition of molasses

a,b,c means with different superscript differ ($p < 0.05$)

Lignin is a part of plant cell walls which consists of very complex and difficult to digest bonds. The high content of lignin in feed will reduce its digestibility. The lignin content of oil palm fronds is 30.18% [2]. One of the purposes of processing feed through ammonization and fermentation in oil palm fronds is to reduce the lignin content. The addition of different additive gave a very significant effect ($P < 0.01$) on the lignin content of FOPF. The lowest lignin content in treatment A ie (19.94%) was not significantly different ($P > 0.05$) compared to treatment C (21.57%) but was very significant ($P < 0.01$) lower treatment B (23.15%) and treatment D (22.41%).

The lowest lignin content in treatment A (19.94%) allegedly in the oil palm frond fermentation using poultry manure will produce higher lignase and urease enzymes than other treatments. The lignase enzyme is an extracellular enzyme that functions to degrade lignin and urease to break down urea into ammonia. Poultry manure also functions as a source of urease which will break down urea into ammonia and CO_2 . The high N content in poultry manure, which is 3.63-5.30% with C/N ratio 5.8-7.6 [16], causes high urease activity so that the addition of poultry manure as a source of additive in palm frond fermentation can reduce lignin content. Thambirajah et al [17] there is a positive

correlation between the nitrogen content of poultry manure and the nutritional content of oil palm empty fruit bunches fermented with poultry manure. The addition of 15% poultry manure in ammoniated rice straw can reduce fermentation time from 20 days to 5 days [18].

Bioactive compounds from extracts of FOPF

Bioactive compounds contained in extracts from FOPF can be known through maceration method with multilevel extraction. This is based on multilevel extraction can separate the bioactive compounds contained in the extract from the FOPF in the same sample based on their level of polarity without the bioactive compounds dissolving in other solvents. Therefore in this study the multilevel extraction process starts from a non-polar solvent (n hexane) followed by a semi-polar solvent (ethyl acetate) and finally with a polar solvent (methanol).

Table 3 shows n acetate extracts FOPF produce the lowest yields ranging from 0.78-1.08% while the methanol extract produces the largest yields ranging from 5.79-7.87%. The same results were reported by Salamah et al [19] the lowest yield of Taiwanese gravestone extract was from n-hexane extract (0.04%) and the highest yield was from methanol extract (0.12%).

Tabel 3. The yield of extracts FOPF

Treatment	Yield of extracts (%)		
	N heksan	Etil Asetat	Metanol
A	0,78	1,53	7,87
B	1,08	1,30	7,07
C	1,02	1,17	5,79
D	0,89	1,08	7,40

Notes : A = FOPF with the addition of poultry manure
 B = FOPF with the addition of cow feces
 C = FOPF with the addition of urea

D = FOPF with the addition of molasses

FOPF which was extracted with methanol, produced the highest yield of 7.87%. This shows that the most bioactive compounds found in FOPF are polar because they can dissolve in polar solvents namely methanol.

Bioactive compounds of FOPF which are non-polar and semipolar in smaller amounts. Phytochemical analysis of bioactive compounds of extract of fermented oil palm fronds is shown in Table 4.

Table 4. Bioactive compounds of extract of FOPF

Class of compounds	n hexane				Etil Asetat				Metanol			
	A	B	C	D	A	B	C	D	A	B	C	D
Alkaloid	-	-	-	-	-	-	-	-	-	-	-	-
Steroid	+	+	+	+	+	+	+	+	+	+	+	+
Terpenoid	-	-	-	-	-	-	-	-	-	-	-	-
Flavanoid	-	-	-	-	-	-	-	-	-	-	-	-
Saponin	-	-	-	-	-	-	-	-	-	-	-	-
Kuinon	-	-	-	-	-	-	-	-	-	-	-	-
Tannin	-	-	-	-	-	-	-	-	+	-	+	-
Kumarin	-	-	-	-	-	-	-	-	-	-	-	-
Fenolik	-	-	-	-	-	-	-	-	+	++	+	++

Notes : A = FOPF with the addition of poultry manure
 B = FOPF with the addition of cow feces
 C = FOPF with the addition of urea
 D = FOPF with the addition of molasses
 - there are no bioactive compounds
 + in small quantities
 ++ there are in sufficient quantities

The extraction process uses methanol solvents in treatments A and C, produce small amounts of tannin and phenolic compounds (+). Sufficient amounts of phenolic compounds (++) were found in treatment B and D. N hexane extract in treatment A produced the lowest yield (0.78%) and methanol extract produced the highest yield (7.87%). This shows that treatment A contains polar compounds (methanol) which are greater than non-polar compounds (n hexane). Analysis of

phytochemical compounds shows, treatment A contains bioactive compounds, are steroids and tannins, in small amounts (+) and phenolic compounds in sufficient quantities (++).

a. Steroid

Steroids are initially considered only as components of animal substances (as sex hormones, adrenal hormones, bile acids, etc.), but recently steroids have also been found in plant substances [20]. Steroids contained in

methanol extract, ethyl acetate and n hexane in treatment A are thought to be adrenal hormones and sex hormones, because in poultry, metabolic waste that cannot be used again is released as excreta through cloaca. Excreta consists of feces, urine and sperm/egg cells. Bose et al [21] stated that steroid compounds were detected in *Achatina fulica* (Freshwater gastropods such as golden snails). Steroids are thought to increase body stamina (aphrodisiac) and anti-inflammation. Silva et al. [22] reported extracts from the leaves of *Agave attenuate*, containing steroid compounds that have anti-inflammatory activity.

Phytochemical analysis showed that bioactive steroid compounds were detected in all extract of FOPF (n hexane, ethyl acetate and methanol) which had different polarity. Harbone [20] states, steroids are cholesterol, which are non-polar so they can dissolve in non-polar solvents, (n hexane), semi-polar (ethyl acetate) and polar (methanol) in small amounts (+). The steroid content in beef, which is extracted using non-polar and polar compounds, shows no difference [23].

Bioactive compounds of steroid, which are found in extract of FOPF can be used as antimicrobials because steroids are bioactive compounds that function as anti-fungal. N hexane extract from lotus seeds contains bioactive steroid compounds, and can inhibit the growth of Gram negative bacteria *Aeromonas hydrophila* and Gram positive bacteria *Streptococcus agalactiae* and *Saprolegnia* sp fungi [24].

b. Tannin

Methanol extract from FOPF in treatments A and C contains small amounts of tannin compounds (+). The tannin compounds found in oil palm plants are reported by Sasidharan [7] in methanol extract of palm oil leaves; Febrina et al [6] on the ethanol extract of oil palm fronds.

The presence of tannin compounds in methanol extract in treatments A and C shows that methanol extract from FOPF has the potential to be antimicrobial. Tannin has antibacterial properties by precipitation proteins because it is suspected that tannins have the same effect as phenolic compounds [25]. The tannin compounds found in the ethanol extracts of oil palm fronds [7] and mangosteen peel [26] have the potential to be antimicrobial because they can inhibit the growth of *S. aureus* and *E. coli*.

c. Phenolic

Phytochemical analysis in treatment B and D showed that sufficient quantities of phenolic bioactive compounds (++) . This indicates that the extract of oil palm frond which is fermented with cow feces and molasses can be used as an antimicrobial. Phenol compounds can be used as antibacterial because they contain aromatic rings with one or two hydroxyl groups that can change the permeability of the cytoplasmic membrane thus inhibiting bacterial growth [20,27]. Ethyl acetate extract of Binahong leaves, containing polyphenols and saponins with the Minimum Inhibitory Concentration (MIC) against *Shigella flexneri* was 8% b/v [28].

Antimicrobial activity test of extracts from FOPF (In Vitro)

Figures 2 a and b show that methanol extract from oil palm fronds which are fermented with different additive has antimicrobial activity which is characterized by the formation of inhibitory zones. The formation of inhibition zones shows that it has inhibited the growth of gram-positive bacteria (*S. aureus*) and gram-negative bacteria (*E. coli*), by compounds contained in methanol extract from FOPF . The inhibitory zone formed shows that bacteria have antimicrobial activity [29], thus it can be concluded that methanol extract from FOPF which are fermented with different sources of additives has antimicrobial activity that is bacteriocide (killing bacteria).

The inhibitory zone formed in testing the antimicrobial activity of extracts from FOPF in Gram positive (*S.aureus*) and Gram negative

(*E.coli*) bacteria at concentrations of 1-5% as shown in Table 5.

Tabel 5. Inhibited Zone Diameter of Extracts from FOPF

Concentration	Inhibited Zone Diameter (mm)							
	<i>E. coli</i>				<i>S. aureus</i>			
	A	B	C	D	A	B	C	D
1 %	12,83	9,00	9,00	7,67	8,00	7,67	11,33	6,30
2 %	11,67	8,67	8,83	7,33	9,16	8,67	10,16	7,67
3 %	10,50	8,66	9,00	7,50	9,67	8,00	10,16	9,16
4 %	8,50	10,33	8,00	8,00	7,83	9,33	8,00	11,67
5 %	7,33	9,00	8,83	7,50	7,83	6,67	9,00	10,00

Notes : A = extracts from FOPF with the addition of poultry manure
 B = extracts from FOPF with the addition of cow feses
 C = extracts from FOPF with the addition of urea
 D = extracts from FOPF with the addition of molasses

Data Table 5 shows the difference in diameter of inhibition zones in each treatment. The diameter of the inhibition zone ranges from 6.67 - 12.83 mm. The smallest diameter inhibition zone of 6.67 mm was produced at a concentration of 5% in treatment B. The largest diameter inhibition zone is 12.83 mm at a concentration of 1% in treatment A.

The ability of extracts from FOPF inhibits bacterial growth in the moderate to strong because the diameter of the inhibition zone ranges from 6.67-12.83 mm. In accordance with the statement of Davis and Stout [30] and Mudi and Ibrahim [31] if the diameter of the inhibition zone is more than 6 mm, the extract has antimicrobial activity with a moderate to strong category.

Table 5 shows the diameter of the inhibitory zone in gram negative bacteria (*Escherichia coli*) generally greater than the gram-positive bacteria (*S. aureus*). The largest diameter of the inhibition zone in gram negative bacteria (*E. coli*) namely in treatment A with a concentration of 1% was 12.83 mm. The largest diameter of the inhibition zone in gram-positive bacteria (*S. aureus*) namely in treatment D with a concentration of 4% is 11.67 mm

There was a difference in the diameter of the inhibitory zone in gram positive and gram negative bacteria due to differences in the structure of bacterial cell walls. The data in Table 5 shows that the antibacterial compounds of the methanol extract of FOPF were able to inhibit the growth of gram negative bacteria better than gram-positive bacteria which was characterized by a larger diameter of the inhibition zone. Cell walls of gram-positive bacteria have a simple structure compared to gram negative bacteria that have complex structures. The largest diameter inhibition zone (12.83 mm) for gram negative bacteria (*E. coli*) showed that the compounds contained in the methanol extract of FOPF had a greater ability to penetrate the cell walls of gram negative bacteria. This is because gram negative bacteria have a more complex structure and have more layers so that only certain compounds can penetrate it.

The Minimum Inhibition Concentration (MIC) shows the lowest concentration of extract which can inhibit bacterial growth. The smaller the MIC value the greater the antimicrobial activity. This can be seen in treatment A at a concentration of 1% which was able to inhibit the growth of pathogenic

microbes with the largest diameter of inhibition zone (12.83 mm).

Conclusion.

FOPF with addition of poultry manure (treatment A), yields the lowest pH value (5.18), sour aroma (score 3), yellowish green (score 3.03) and softer texture (score 2) and the lowest lignin content (19.94%). Methanol extract from FOPF with addition of poultry manure (treatment A) produces the highest extract of rendemen (7.87%) and the largest inhibition zone diameter (12.83 mm) with MIC is 1%.

Acknowledgment.

Author would like to thank to the Institute for Research and Community Service of the State Islamic University of Sultan Syarif Kasim Riau which funded this study under contract number 1043/R/2017.

References

1. Febrina, D., N. Jamarun, M. Zain and Khasrad. The effects of P, S and Mg supplementation of oil palm fronds fermented by *Phanerochaete chrysosporium* on rumen fluid characteristics and microbial protein synthesis. *Pakistan Journal of Nutrition*. 2016; 15:299-304.
2. Febrina, D., N. Jamarun, M. Zain and Khasrad. Effects of calcium (Ca) and manganese (Mn) supplementation during oil palm frond fermentation by *Phanerochaete chrysosporium* on in vitro digestibility and rumen fluid characteristics. *Pakistan Journal of Nutrition*. 2016; 15: 352-358.
3. Zahari W. M. ., O. A. Hassan., H. K. Wong and J. B. Liang. Utilization of Oil Palm Frond - Based Diets for Beef and Dairy Production in Malaysia. *Asian-Aust. J. Anim. Sci.* 2003; 16 (4):625-634.
4. Rahman. M.M., M. Lourenço., H.A. Hassim., J.J.P. Baars., A.S.M. Sonnenberg., J.W. Cone., J.De Boever., V.Fievez. Improving ruminal degradability of oil palm fronds using white rot fungi. *Animal Feed Science and Technology*. 2011; 169. Issues 3–4:157-166.
5. Kawamoto, H., Mohamed, W.Z., Sukur, N.I.M., Ali, M.S.M., Islam, Y., Oshio, S. Palatability, digestibility and voluntary intake of processed oil palm fronds in cattle. *Japan Agric. Res. Quart.* 2001; 35,195–200.
6. Febrina. D., R. Febriyanti., S. I. Zam., J. Handoko., A. Fatah and J. Juliantoni. Antibacterial Activity Testing and Ethanol Extract Characterization of Oil Palm Fronds (*Elaeis guineensis* Jacq). *Pakistan Journal of Nutrition*. 2018; 17 (9): 427-433.
7. Sasidharan. S., Rajoo. N. Rathinam. X., Lachimanan. Y.L., and Rajoo. A. Wound Healing Potential of *Elaeis guineensis* Jacq Leaves in an Infected Albino Rat Model. *Molecules*. 2010; 15:3186-3199.
8. Syahmi, A.R.M.; Vijayarathna, S.; Sasidharan, S.; Latha, L.Y.; Kwan, Y.P.; Lau, Y.L.; Shin, L.N.; Chen, Y. Acute oral toxicity and brine shrimp lethality of *Elaeis guineensis* jacq., (oil palm leaf) methanol extract. *Molecules*. 2010; 15, 8111–8121.
9. Febrina, D. Utilization of oil palm fronds biodelignification products using *Phanerochaete chrysosporium* as substitute forage on goat. Ph.D. Thesis,

- Faculty of Animal Science, University of Andalas, Padang, Indonesia. 2016.
10. Franswort, N. R. Biological and Phytochemical Screenings of Plant. *J. Pharm. Sci.*, 1996; 55 (3): 225-265.
 11. Handayani, D., N. Sandrawaty, M. Murniati and R. Regina. Screening of endophytic bacteria isolated from marine sponge *Haliclona fascigera* for inhibition against clinical isolates of methicillin resistant *Staphylococcus aureus* (MRSA). *J. Applied Pharmaceut. Sci.* 2015; 5: 139-142.
 12. Ozduven. M. L., F. KOC., C. Polat., L. Coskuntuna. The Effects of Lactic Acid Bacteria and Enzyme Mixture Inoculants on Fermentation and Nutrient Digestibility of Sunflower Silage. *Kafkas Univ Vet Fak Derg* 2009; 15 (2): 195-199.
 13. Laura. B and G. Paolo. Algae. Anatomy Biochemistry and Biotechnology. CRC Press. Boca Raton. New York. 2006.
 14. Umiyasih. U dan E. Wina. Processing and Nutritional Value of Corn by-Product as Ruminant Feed. *Wartazoa.* 2008; 18(3):127-136.
 15. Saun, R.J.V, & A.J. Henrich. Trouble shooting silage problem: How to identify potential problem. In: Proceedings of the Mid-Atlantic Conference: Pennsylvania, 26 May 2008. Penn State's Collage. 2008. pp 2-10.
 16. Amanullah. M. A., S. Sekar and P. Muthukrishnan. Prospect and Potential of Poultry Manure. *Asian Journal of Plant Sciences.* 2010; 1 – 11.
 17. Thambirajah. J. J., M.D.Zulkali., M.A.Hashim. Microbiological and biochemical changes during the composting of oil palm empty-fruit-bunches. Effect of nitrogen supplementation on the substrate. *Bioresource Technology.* 1995; 52(2): 133-144.
 18. Salamah E., E. Ayuningrat dan S. Purwaningsih S. Initial Screening of Fresh Water Mussel Bioactive Components from (*Anodonta woodiana* Lea.) from Antioxidant Compounds. *Buletin Teknologi Hasil Perikanan.* 2008; 11(2):119-132.
 19. Harborne.A. J. Phytochemical Methods A Guide to Modern Techniques of Plant Analysis. Edition Number 3. Springer Netherlands Number of Pages XIV, 302. 1998.
 20. Bose R, Majumdar C, Bhattacharya S. Steroids in *Achatina fulica* (Bowdich): steroid profile in haemolymph and in vitro release of steroids from endogenous precursors by ovotestis and albumen gland. *Comp Biochem Physiol.* 1997; 116C(3):179-182.
 21. Silva BP, Sousa AC, Silva GM, Mendes TP, Parente JP. A new bioactive steroidal saponin from *Agave attenuata*. *Z Naturforsch* 2002; 57C:423-428.
 22. Schmidt G and H. Steinhart H. Impact of extraction solvents on steroid contents determined in beef. *Journal of Food Chemistry.* 2001; 76:83-88.
 23. Widya. D. R., D. Suryanto dan Desrita. Antimicrobial Activity of Lotus Seeds (*Nymphaea pubescens* L.) on Bacteria *Aeromonas hydrophila*, *Streptococcus agalactiae*, and *Saprolegnia* sp. *Jurnal Aquacoastmarine.* 2104; 2(1).7-14.

24. Martono. B dan R. T. Setiyono. Phytochemical Screening of Six Tea Genotypes J. TIDP 2014; 1(2), 63-68.
25. Poelongan and Praptiwi. Antibacterial Activity Test of Mangos Teen (*Gardnia mangostana* Linn) Peel. Media Litbang Kesehatan. 2010; 20 (2):65-69.
26. Ikalinus. R., S. K. Widyastuti dan Ni. L. E. Setiasih. Phytochemical Screening Ethanol Extract Skin Stem Moringa (*Moringa oleifera*). Indonesia Medicus Veterinus. 2015; 4(1):71-79.
27. Wardhani, L. K. dan N. Sulistyani. Antibacterial Activity Test of Ethyl Acetate Extract of Binahong Leaf (*Anredera scandens* (L.) Moq.) Against *Shigella flexneri* with The Profile of Thin Layer Chromatography. Jurnal Ilmiah Kefarmasian. 2012; 2(1):1-16.
28. Nanda. A., M. and Saravanan. Biosynthesis of silver nanoparticles. from *Staphylococcus aureus* and its antimicrobial activity against MRSA and MRSE. Nanomedicine: Nanotechnology, Biology and Medicine. 2009; 5(4): 452-456.
29. Davis, W.W. and T.R. Stout, Disc plate method of microbiological antibiotic assay. I. Factors influencing variability and error. Applied Microbiol. 1971; 22: 659-665.
30. Mudi, S.Y. and H. Ibrahim, Activity of *Bryophyllum pinnatum* S. Kurz extracts on respiratory tract pathogenic bacteria. Bayero J. Pure Applied Sci. 2008; 1: 43-48.

PEMBERITAHUAN DARI EDITOR

“MAJOR REVISION”

15:54

4G+ 46%



[JTLS] Editor Decision Inbox



Nashi Widodo 10/10/2019

to me ▾



Mr Syukria Ikhsan Zam:

We have reached a decision regarding your submission to Journal of Tropical Life Science, "Nutritional Content and Characteristics of Antimicrobial Compounds from Fermented Oil Palm Fronds (*Elaeis guineensis* Jacq.)".

Our decision is to: MAJOR REVISION. Please effect the corrections as soon as possible.

Prof. Dr. Fahrul zaman Huyop
Universiti Teknologi Malaysia
Phone 0126849374
Fax 0126849374
fahrul@utm.my

Fahrul Huyop
Journal of Tropical Life Science
Universitas Brawijaya

HASIL TELAAH REVIEWER

ABSTRACT

This research was conducted to determine the nutritional quality and characteristics of antimicrobial compounds from fermented oil palm fronds (FOPF). This research was carried out using a Complete Randomized Design with 4 treatments and 4 replications. The following treatments were performed by the addition of different additives namely: 10% poultry manure, 10% cow feces, 5% urea and 5% molasses. The data were analyzed using a one-way analysis of variance and differences between treatments were tested using Duncan's multiple range test. The observed parameters were physical quality tests include pH, aroma, color, texture and nutritional quality including the content of NDF- [write in full do not use abbreviations](#), ADF, hemicellulose, cellulose and lignin. Extraction of oil palm fronds using multilevel maceration method with ethyl acetate, n-hexane and methanol. Identification [of what?](#) is done through phytochemical screening. The activity test of antimicrobial compound extracts from FOPF using paper disc method with a concentration of 1-5. FOPF with poultry manure yields the lowest pH value, sour aroma, yellowish green, softer texture and the lowest lignin content (19.94%). Methanol extract from FOPF with poultry manure produces the highest extract yield and the largest inhibition zone diameter with a MIC is 1%.

[Please](#) [add](#) [in](#) [conclusion](#) [of](#) [studies?](#)

Background

The utilization of oil palm plantation waste such as oil palm fronds as feed is still limited due to the high lignin content [1, 2]. Physical-chemical-biological and combination treatments and the addition of additives on the fermentation process aim to break down lignocellulose and ligno hemicellulose bonds and increase nutrient digestibility [3, 4, 5].

Palm oil fronds have antimicrobial activity [6]. Methanol extract from palm oil leaves contains tannin, alkaloids, saponins, terpenoids and flavonoids [7]; and toxicity tests on shrimp and rats showed that palm oil leaf extract is safer for consumption and recommended as a natural product [8]. This study aims to increase the nutritional content of oil palm fronds through a fermentation process with the addition of different additives and to determine the bioactive compounds in extracts from FOPF with multilevel extraction using n hexane, ethyl acetate and methanol.

Methods

The research was conducted in June-November 2017, at the Laboratory of Nutrition and Chemistry at the Faculty of Agriculture and Animal Sciences of State Islamic University of Sultan Syarif Kasim Riau Indonesia, Biotechnology Research Center of the Indonesian Sciences Institute Indonesia and

Laboratory of Biota Sumatra Andalas University.

Materials

The materials used in this research are NDS- [write in full?](#) solution, ADS- [write in full?](#) solution, acetone, alcohol, octanol, H₂SO₄, methanol, ethyl acetate, hexane, filter paper, rotary vacuum evaporator, HCl, ammonia, Mayer reagent, Dragendorff reagent, ether, Liebermann Bouchard reagent, Mg powder, amyl alcohol, NaOH, FeCl₃, aquadest, poultry manure, cow feces, urea, molasses, plastic bags. The tools used in the study were leaf chopper, ~~eruisel dish, fibertec hot extraction, fibertec cold extraction, oven, desicator, furnace, analytical scales, dropple plate, water bath.~~

Research Procedure

The study was conducted through two (2) stages: namely fermentation of oil palm frond based on Febrina [9] modified with different additive (10% poultry manure, 10% cow feces, 5% urea and 5% molasses) using a Completely Randomized Design 4 treatment with 4 replications. Fermentation process for 21 days. The data were analyzed using a one-way analysis of variance and differences between treatments were tested using Duncan's Multiple Range Test. The observed parameters were physical quality tests include pH, aroma, color and texture and nutritional quality including the

Commented [1]: Pls delete

content of NDF, ADF, hemi cellulose, cellulose and lignin based on Van Soest analysis using the FOSS Fibertec system. The second stage is multilevel maceration and extraction using methanol, ethyl acetate and n-hexane solvents. Phytochemical screening based on Franswort [10] and the antibacterial activity test of oil palm fronds extract based on Handayani *et al.* [11].

Result and Discussion

The Effect of Different Additives on The Physical quality of FOPF

The addition of different additives has a significant effect ($P < 0.05$) on the physical quality of FOPF including pH, aroma, color and texture (Table 1).

Table 1. The effect of different additives on the physical quality of FOPF

Treatment	Physical quality							
	pH		Aroma		Color		Texture	
	Value	Notes	Value	Notes	Value	Notes	Value	Notes
A	5,18 ^a ±0,10	Acid	3,00 ^b ±0,00	Acid	3,03 ^a ±0,05	Yellowish Green	2,00 ^a ±0,00	Rather Soft
B	5,52 ^b ±0,20	Acid	4,00 ^c ±0,00	More Acidic	3,00 ^a ±0,00	Yellowish Green	3,00 ^b ±0,00	Soft
C	7,60 ^c ±0,14	Base	2,03 ^a ±0,05	Non-acidic	4,03 ^b ±0,05	Brownish Green	3,03 ^b ±0,05	Soft
D	5,28 ^{ab} ±0,21	Acid	5,00 ^d ±0,00	Very Acidic	3,00 ^a ±0,00	Yellowish Green	3,00 ^b ±0,00	Soft

Notes: A = FOPF with the addition of poultry manure
 B = FOPF with the addition of cow feces
 C = FOPF with the addition of urea
 D = FOPF with the addition of molasses
 a,b,c means with different superscript differ ($p < 0.05$)

The lowest of pH (5.18) in treatment A is related to the availability of carbohydrates and the production of lactic acid bacteria. The same results were seen in treatment D with a pH of 5.28. Addition of poultry manure and molasses on fermentation oil palm frond aims as a source of additive which can trigger the growth of lactic acid bacteria to produce lactic acid. Increased production of lactic acid will reduce pH and inhibit the growth of pathogenic bacteria. Ozduven *et al.* [12] stated that pH is associated with lactic acid produced during the fermentation process which can inhibit the growth of pathogenic microbes. Oil palm fronds which are fermented with molasses produce pH 3.93 [3].

The process of ammonization of oil palm fronds using urea will produces the highest pH of 7.60

which is alkaline. The same results were reported by Zahari *et al.* [3], the process of ammonization of oil palm fronds using urea resulting in a pH of 7.38. In the process of ammonization with urea will break down into ammonium ions so that the ammonization product is alkaline Laura and Paolo [13] state that urea will break down into ammonium ions, which are alkaline and ammonization processes can break down glycoside bonds [14] so that the oil palm fronds can be degraded by rumen microbes.

In treatment A produces an

aroma score of 3 with acid criteria. This shows that the fermentation process is good which is characterized by acidic products were typical of

Commented [2]: Please use past tense in all sentences. Is become was etc.

fermentation. Aroma is related to the production of lactic acid and pH it can be seen in Table 1 the addition of poultry manure to oil palm fronds fermentation resulting in the lowest pH (5.18) so that the products are acidic.

Treatment A, B and D produces a color that is not different ($P > 0.05$), which is natural green or yellowish green with score of 3-3.03. Good quality silage will produce colors that are not different from the original material [15] The occurrence of color changes from green to yellowish green indicated due to the anaerobic phase oxygen has run out of then carbohydrates. Then it will produce CO₂, H₂O and heat. The heat produced causes a change in color from green to yellowish green.

Treatment A will produce softer textures with score 2. This shows that the addition of the

poultry manure to the fermentation process can support the growth of lactic acid bacteria to produce lactic acid. Optimal production of lactic acid will reduce pH so that it inhibits the growth of pathogenic bacteria. This low pH also supports the production of lignase enzymes which function to break down lignocellulose and ligno-hemicellulose thereby reducing lignin content. It is seen that treatment A produced the lowest lignin content of 19.94%. The release of lignin from cellulose and hemicellulose causes fermented products to have a softer texture.

The Effect of Different Additive on Fiber Fraction of FOPF

The addition of different additive had no effect ($P>0.05$) on the content of NDF, ADF, hemicellulose and cellulose but had a very significant effect ($P<0.01$) on the lignin content (Table 2). Rahman *et al.* [4] reported there was a decrease in the lignin content of oil palm fronds in the fermentation process using white rot fungi.

The lowest lignin content in treatment A (19.94%) allegedly in the oil palm frond fermentation using poultry manure will produce higher lignase and urease enzymes than other treatments. The lignase enzyme is an extracellular enzyme that functions to degrade lignin and urease to break down urea into ammonia. Poultry manure also functions as a source of urease which will break down urea into ammonia and CO₂. The high N content in poultry manure, which is 3.63-5.30% with C/N ratio 5.8-7.6 [16], causes high urease activity so that the addition of poultry manure as a source of additive in palm frond fermentation can reduce lignin content. Thambirajah *et al.* [17] there is a positive correlation between the nitrogen content of poultry manure and the nutritional content of oil palm empty fruit bunches fermented with poultry manure. The addition of 15% poultry manure in ammoniated rice straw can reduce fermentation time from 20 days to 5 days.

Table 2. The effect of different additive on fiber fraction of FOPF.

Treatment	NDF	ADF	Fiber Fraction (%)		
			Hemicellulose	Cellulose	Lignin
A	73,34±3,68	51,81±4,82	21,52±6,07	28,54±3,36	19,94 ^a ±1,47
B	75,39±2,03	55,03±4,23	20,35±4,46	29,57±3,09	23,15 ^b ±0,96
C	73,83±1,01	53,30±3,43	20,51±4,30	31,55±1,99	21,57 ^a ±1,58
D	74,98±3,52	54,04±4,48	20,93±6,63	31,70±4,27	22,41 ^b ±1,22

Notes: A = FOPF with the addition of poultry manure
 B = FOPF with the addition of cow feces
 C = FOPF with the addition of urea
 D = FOPF with the addition of molasses

Lignin is a part of plant cell walls which consists of very complex and difficult to digest bonds. The high content of lignin in feed will reduce its digestibility. The lignin content of oil palm fronds is 30.18% [2]. One of the purposes of processing feed through ammonization and fermentation in oil palm fronds is to reduce the lignin content. The addition of different additive gave a very significant effect ($P < 0.01$) on the lignin content of FOPF. The lowest lignin content in treatment A (19.94%) was not significantly different ($P > 0.05$) compared to treatment C (21.57%) but was very significant ($P < 0.01$) lower treatment B (23.15%) and treatment D (22.41%).

Bioactive Compounds from Extracts of FOPF

Bioactive compounds contained in extracts from FOPF can be known through maceration method with multilevel extraction. This is based on multilevel extraction can separate the bioactive compounds contained in the extract from the FOPF in the same sample based on their level of polarity without the bioactive compounds dissolving in other solvents, therefore in this study the multilevel extraction process starts from a non-polar solvent (n-hexane) followed by a semi-polar solvent (ethyl acetate) and finally with a polar solvent (methanol).

Table 3 shows n acetate extracts FOPF produce the lowest yields ranging from 0.78-1.08% while

the methanol extract produces the largest yields ranging from 5.79-7.87%. The same results were reported by Salamah *et al.* [18] the lowest yield of Taiwanese gravestone extract was from n-hexane extract (0.04%) and the highest yield was from methanol extract (0.12%).

FOPF which was extracted with methanol, produced the highest yield of 7.87% [of what compound?](#). This shows

Table 3. The yield of extracts FOPF
Yield of extracts (%)

Treatment	N heksan	Etil	Metanol	Asetat		
				A	B	C
A	0,78	1,53	7,87			
B	1,08	1,30	7,07			
C	1,02	1,17	5,79			
D	0,89	1,08	7,40			

Notes: A = FOPF with the addition of poultry manure
B = FOPF with the addition of cow feces
C = FOPF with the addition of urea
D = FOPF with the addition of molasses

Table 4. Bioactive compounds of extract of FOPF

Class of Compounds	n hexane			Ethyl Acetate			Methanol			
	A	B	C	A	B	C	A	B	C	D
Alkaloid	-	-	-	-	-	-	-	-	-	-
Steroid	+	+	+	+	+	+	+	+	+	+
Terpenoid	-	-	-	-	-	-	-	-	-	-
Flavonoid	-	-	-	-	-	-	-	-	-	-
Saponin	-	-	-	-	-	-	-	-	-	-
Kuinon	-	-	-	-	-	-	-	-	-	-
Tannin	-	-	-	-	-	-	+	-	+	-
Kumarin	-	-	-	-	-	-	-	-	-	-
Fenolik	-	-	-	-	-	-	+	++	+	++

Notes: A = FOPF with the addition of poultry manure
B = FOPF with the addition of cow feces
C = FOPF with the addition of urea
D = FOPF with the addition of molasses
- = there are no bioactive compounds
+ = in small quantities
++ = there are in sufficient quantities

that the most bioactive compounds found in FOPF are polar because they can dissolve in polar solvents namely methanol. Bioactive compounds of FOPF which are non-polar and semi-polar in smaller amounts. Phytochemical

analysis of bioactive compounds of extract of fermented oil palm fronds is shown in Table 4.

The extraction process uses methanol solvents in treatments A and C, produce small amounts of tannin and phenolic compounds (+). Sufficient amounts of phenolic compounds (++) were found in treatment B and D. N hexane extract in treatment A produced the lowest yield (0.78%) and methanol extract produced the highest yield (7.87%). This shows that treatment A contains polar compounds (methanol) which are greater

than non-polar compounds (n hexane). Analysis

of phytochemical compounds shows, treatment A contains bioactive compounds, are steroids and tannins, in small amounts (+) and phenolic compounds in sufficient quantities (++).

a. Steroid
Steroids are initially considered only as

components of animal substances (as sex hormones, adrenal hormones, bile acids, etc.), but recently steroids have also been found in plant substances [19]. Steroids contained in methanol extract, ethyl acetate and n hexane in treatment A are thought to be adrenal hormones and sex hormones, because in poultry, metabolic

waste that cannot be used again is released as excreta through cloaca. Excreta consists of feces, urine and sperm/egg cells. Bose *et al.* [20] stated that steroid compounds were detected in *Achatina fulica* (freshwater gastropods such as golden snails). Steroids are thought to increase

body stamina (aphrodisiac) and anti-inflammation. Silva *et al.* [21] reported extracts from the leaves of *Agave attenuate*, containing steroid compounds that have anti-inflammatory activity.

Phytochemical analysis showed that bioactive steroid compounds were detected in all extract of FOPF (n hexane, ethyl acetate and methanol) which had different polarity. Harbone [19] states, steroids are cholesterol, which are non-

properties by precipitation proteins because it is suspected that tannins have the same effect as phenolic compounds [24]. The tannin compounds found in the ethanol extracts of oil palm fronds [7] and mangosteen peel [25] have the potential to be antimicrobial because they can inhibit the growth of *S. aureus* and *E. coli*.

c. Phenolic

Phytochemical analysis in treatment B and D showed that sufficient quantities of phenolic

Table 5. Inhibited Zone Diameter of Extracts from FOPF

Concentration	Inhibited Zone (mm)							
	<i>E. coli</i>				<i>S. aureus</i>			
	A	B	C	D	A	B	C	D
1 %	12,83	9,00	9,00	7,67	8,00	7,67	11,33	6,30
2 %	11,67	8,67	8,83	7,33	9,16	8,67	10,16	7,67
3 %	10,50	8,66	9,00	7,50	9,67	8,00	10,16	9,16
4 %	8,50	10,33	8,00	8,00	7,83	9,33	8,00	11,67
5 %	7,33	9,00	8,83	7,50	7,83	6,67	9,00	10,00

Notes: A = extracts from FOPF with the addition of poultry manure
 B = extracts from FOPF with the addition of cow feces
 C = extracts from FOPF with the addition of urea
 D = extracts from FOPF with the addition of molasses

polar so they can dissolve in non-polar solvents, (n-hexane), semi-polar (ethyl acetate) and polar (methanol) in small amounts (+). The steroid content in beef, which is extracted using non-polar and polar compounds, shows no difference [22].

Bioactive compounds of steroid, which are found in extract of FOPF can be used as antimicrobials because steroids are bioactive compounds that function as anti-fungal. N hexane extract from lotus seeds contains bioactive steroid compounds and can inhibit the growth of Gram-negative bacteria (*Aeromonas hydrophila*) and Gram-positive bacteria (*Streptococcus agalactiae*) and fungi (*Saprolegnia* sp.) [23].

b. Tannin

Methanol extract from FOPF in treatments A and C contains small amounts of tannin compounds (+). The tannin compounds found in oil palm plants are reported by Sasidharan [7] in methanol extract of palm oil leaves; Febrina *et al.* [6] on the ethanol extract of oil palm fronds. The presence of tannin compounds in methanol extract in treatments A and C shows that methanol extract from FOPF has the potential to be antimicrobial. Tannin has antibacterial

bioactive compounds (++). This indicates that the extract of oil palm frond which is fermented with cow feces and molasses can be used as an antimicrobial. Phenol compounds can be used as antibacterial because they contain aromatic rings with one or two hydroxyl groups that can change the permeability of the cytoplasmic membrane thus inhibiting bacterial growth [19,26]. Ethyl acetate extract of binahong leaves, containing polyphenols and saponins with the minimum inhibitory concentration (MIC) against *Shigella flexneri* was 8% b/v [27].

Antimicrobial activity test of extracts from FOPF (In Vitro)

Figures 2 a and b show that methanol extract from oil palm fronds which are fermented with different additive has antimicrobial activity which is characterized by the formation of inhibitory zones. The formation of inhibition zones shows that it has inhibited the growth of Gram-positive bacteria (*S. aureus*) and Gram-negative bacteria (*E. coli*), by compounds contained in methanol extract from FOPF. The inhibitory zone formed shows that bacteria have antimicrobial activity [28], thus it can be concluded that methanol extract from FOPF

which are fermented with different sources of additives has antimicrobial activity that is bactericide (killing bacteria).

The inhibitory zone formed in testing the antimicrobial activity of extracts from FOPF in Gram-positive (*S. aureus*) and Gram-negative (*E. coli*) bacteria at concentrations of 1-5% as shown in Table 5.

~~In Data-~~ Table 5 shows the difference in diameter of inhibition zones in each treatment. The diameter of the inhibition zone ranges from 6.67 - 12.83 mm. The smallest diameter inhibition zone of 6.67 mm was produced at a concentration of 5% in treatment B. The largest diameter inhibition zone is 12.83 mm at a concentration of 1% in treatment A.

The ability of extracts from FOPF inhibits bacterial growth in the moderate to strong, because the diameter of the inhibition zone ranges from 6.67-12.83 mm. In accordance with the statement of Davis and Stout [29] and Mudi and Ibrahim [30] if the diameter of the inhibition zone is more than 6 mm, the extract has antimicrobial activity with a moderate to strong category.

Table 5 shows the diameter of the inhibitory zone in Gram-negative bacteria (*E. coli*) generally greater than the Gram-positive bacteria (*S. aureus*). The largest diameter of the inhibition zone in Gram-negative bacteria (*E. coli*) namely in treatment A with a concentration of 1% was 12.83 mm. The largest diameter of the inhibition zone in Gram-positive bacteria (*S. aureus*) namely in treatment D with a concentration of 4% is 11.67 mm?? No full stop?

There was a difference in the diameter of the inhibitory zone in gram positive and Gram-negative bacteria due to differences in the structure of bacterial cell walls. The data in Table 5 shows that the antibacterial compounds of the methanol extract of FOPF were able to inhibit the growth of Gram-negative bacteria better than Gram-positive bacteria which was characterized by a larger diameter of the inhibition zone. Cell walls of Gram-positive bacteria have a simple structure compared to Gram-negative bacteria that have complex structures. The largest diameter inhibition zone (12.83 mm) for gram negative bacteria (*E. coli*) showed that the compounds contained in the

methanol extract of FOPF had a greater ability to penetrate the cell walls of Gram-negative bacteria. This is because Gram-negative bacteria have a more complex structure and have more layers so that only certain compounds can penetrate it.

The minimum inhibition concentration (MIC) shows the lowest concentration of extract which can inhibit bacterial growth. The smaller the MIC value the greater the antimicrobial activity. This can be seen in treatment A at a concentration of 1% which was able to inhibit the growth of pathogenic microbes with the largest diameter of inhibition zone (12.83 mm).

Conclusion

FOPF with addition of poultry manure (treatment A), yields the lowest pH value (5.18), sour aroma (score 3), yellowish green (score 3.03) and softer texture (score 2) and the lowest lignin content (19.94%). Methanol extract from FOPF with addition of poultry manure (treatment A) produces the highest extract of rendemen (7.87%) and the largest inhibition zone diameter (12.83 mm) with MIC is 1%.

[So this will suggests: Revise conclusion to show benefits of the study?](#)

Acknowledgement

Author would like to thank to the Institute for Research and Community Service of the State Islamic University of Sultan Syarif Kasim Riau which funded this study under contract number 1043/R/2017.

References

1. Febrina D, Jamarun N, Zain M, Khasrad (2016) The effects of P, S and Mg supplementation of oil palm fronds fermented by *Phanerochaete chrysosporium* on rumen fluid characteristics and microbial protein synthesis. Pakistan Journal of Nutrition 15: 299 – 304. doi: 10.3923/pjn.2016.299.304.
2. Febrina D, Jamarun N, Zain M, Khasrad (2016) Effects of calcium (Ca) and manganese (Mn) supplementation during oil palm frond fermentation by *Phanerochaete chrysosporium* on in vitro digestibility and rumen fluid characteristics.

- Pakistan Journal of Nutrition 15: 352 – 358. doi: 10.3923/pjn.2016.352.358.
3. Zahari WM, Hassan OA, Wong HK, Liang JB (2003) Utilization of oil palm frond - based diets for beef and dairy production in Malaysia. *Asian-Australasian Journal of Animal Sciences* 16 (4): 625 – 634. doi: [10.5713/ajas.2003.625](https://doi.org/10.5713/ajas.2003.625).
 4. Rahman MM, Lourenço M, Hassim HA, Baars JJP, Sonnenberg ASM, Cone JW, de Boever J, Fievez V (2011) Improving ruminal degradability of oil palm fronds using white rot fungi. *Animal Feed Science and Technology* 169 (3-4): 157 – 166. doi: 10.1016/j.anifeedsci.2011.06.014.
 5. Kawamoto H, Mohamed WZ, Sukur NIM, Ali MSM, Islam Y, Oshio S (2001) Palatability, digestibility and voluntary intake of processed oil palm fronds in cattle. *Japan Agricultural Research Quarterly* 35; 195 – 200. doi: [10.6090/jarq.35.195](https://doi.org/10.6090/jarq.35.195).
 6. Febrina D, Febriyanti R, Zam SI, Handoko J, Fatah A, Juliantoni J (2018) Antibacterial activity testing and ethanol extract characterization of oil palm fronds (*Elaeis guineensis* Jacq.). *Pakistan Journal of Nutrition* 17 (9): 427 – 433. doi: [10.3923/pjn.2018.427.433](https://doi.org/10.3923/pjn.2018.427.433).
 7. Sasidharan S, Rajoo N, Rathinam X, Lachimanan YL, Rajoo A (2010) Wound healing potential of *Elaeis guineensis* Jacq. leaves in an infected albino rat model. *Molecules* 15: 3186 – 3199. doi: 10.3390/molecules15053186.
 8. Syahmi ARM, Vijayarathna S, Sasidharan S, Latha LY, Kwan YP, Lau YL, Shin LN, Chen Y (2010) Acute oral toxicity and brine shrimp lethality of *Elaeis guineensis* Jacq. (oil palm leaf) methanol extract. *Molecules* 15: 8111 – 8121. doi: 10.3390/molecules15118111.
 9. Febrina D (2016) Utilization of oil palm fronds biodelignification products using *Phanerochaete chrysosporium* as substitute forage on goat. PhD thesis. Andalas University, Department Animal Science.
 10. Franswort NR (1996) Biological and phytochemical screenings of plant. *Journal of Pharmaceutical Science* 55 (3): 225 – 265. doi: 10.1002/jps.2600550302.
 11. Handayani D, Sandrawaty N, Murniati M, Regina R (2015) Screening of endophytic bacteria isolated from marine sponge *Haliclona fascigera* for inhibition against clinical isolates of methicillin resistant *Staphylococcus aureus* (MRSA). *Journal of Applied Pharmaceutical Science* 5: 139 – 142. doi: 10.7324/JAPS.2015.50926.
 12. Ozduven ML, Koc F, Polat C, Coskuntuna L (2009) The effects of lactic acid bacteria and enzyme mixture inoculants on fermentation and nutrient digestibility of sunflower silage. *Kafkas Univ Vet Fak Derg* 15 (2): 195 – 199. doi: 10.9775/kvfd.2008.91-A.
 13. Laura B, Paolo G (2006) *Algae: Anatomy Biochemistry and Biotechnology*. Boca Raton, CRC Press.
 14. Umiyasih U, Wina E (2008) Processing and nutritional value of corn by-product as ruminant feed. *Wartazoa* 18 (3): 127 – 136.
 15. Saun RJV, Henrich AJ (Eds) (2008) *Proceedings of the Mid-Atlantic Conference: 26 May 2008; Pennsylvania*. Penn States Collage.
 16. Amanullah MA, Sekar S, Muthukrishnan P (2010) Prospect and potential of poultry manure. *Asian Journal of Plant Sciences* 9 (4): 1 – 11. doi: 10.3923/ajps.2010.172.182.
 17. Thambirajah JJ, Zulkali MD, Hashim MA (1995) Microbiological and biochemical

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

- changes during the composting of oil palm empty-fruit-bunches. Effect of nitrogen supplementation on the substrate. *Bioresource Technology* 52 (2): 133 – 144. doi: [10.1016/0960-8524\(95\)00008-3](https://doi.org/10.1016/0960-8524(95)00008-3).
18. Salamah E, Ayuningrat E, Purwaningsih S (2008) Initial screening of freshwater mussel bioactive components from (*Anodonta woodiana* Lea.) from antioxidant compounds. *Buletin Teknologi Hasil Perikanan* 11 (2): 119 – 132. doi: [10.17844/jphpi.v11i2.912](https://doi.org/10.17844/jphpi.v11i2.912).
19. Harborne AJ. *Phytochemical Methods A Guide to Modern Techniques of Plant Analysis*. 3rd Edition. Dordrecht, Springer.
20. Bose R, Majumdar C, Bhattacharya S (1997) Steroids in *Achatina fulica* (bowdich): Steroid profile in haemolymph and in vitro release of steroids from endogenous precursors by ootestis and albumen gland. *Comparative Biochemistry and Physiology* 116 (3): 179 – 182. doi: [10.1016/S0742-8413\(96\)00163-6](https://doi.org/10.1016/S0742-8413(96)00163-6).
21. Silva BP, Sousa AC, Silva GM, Mendes TP, Parente JP. A new bioactive steroidal saponin from *Agave attenuata*. *Zeitschrift für Naturforschung* 57c: 423 – 428.
22. Schmidt G, Steinhart H (2001) Impact of extraction solvents on steroid contents determined in beef. *Food Chemistry* 76 (1): 83 – 88. doi: [10.1016/S0308-8146\(01\)00237-0](https://doi.org/10.1016/S0308-8146(01)00237-0).
23. Widya DR, Suryanto D, Desrita (2014) Antimicrobial activity of lotus seeds (*Nymphaea pubescens* L.) on bacteria *Aeromonas hydrophila*, *Streptococcus agalactiae*, and *Saprolegnia* sp. *Jurnal Aquacoastmarine* 2 (1): 7 – 14.
24. Martono B, Setiyono RT (2014) Phytochemical screening of six tea genotypes. *Journal of Industrial and Beverage Crops* 1 (2): 63 – 68. doi: [10.21082/jtidp.v1n2.2014.p63-68](https://doi.org/10.21082/jtidp.v1n2.2014.p63-68).
25. Poelongan M, Praptiwi (2010) Antibacterial activity test of mangos teen (*Gardnia mangostana* Linn.) peel. *Media Litbang Kesehatan*. 20 (2): 65 – 69.
26. Ikalinus R, Widyastuti SK, Setiasih NLE (2015) Phytochemical screening ethanol extract skin stem moringa (*Moringa oleifera*). *Indonesia Medicus Veterinus* 4 (1): 71 – 79.
27. Wardhani LK, Sulistyani N (2012) Antibacterial activity test of ethyl acetate extract of binahong leaf (*Anredera scandens* (L.) Moq.) against *Shigella flexneri* with the profile of thin layer chromatography. *Jurnal Ilmiah Kefarmasian* 2 (1): 1 – 16.
28. Nanda. A, Saravanan M (2009) Biosynthesis of silver nanoparticles from *Staphylococcus aureus* and its antimicrobial activity against MRSA and MRSE. *Nanomedicine: Nanotechnology, Biology and Medicine* 5 (4): 452 – 456. doi: [10.1016/j.nano.2009.01.012](https://doi.org/10.1016/j.nano.2009.01.012).
29. Davis WW, Stout TR (1971) Disc plate method of microbiological antibiotic assay. I. Factors influencing variability and error. *Applied Microbiology* 22: 659 – 665.
30. Mudi SY, Ibrahim H (2008) Activity of *Bryophyllum pinnatum* S. Kurz extracts on respiratory tract pathogenic bacteria. *Bayero Journal of Pure Applied Sciences* 1 (1): 43 – 48. doi: [10.4314/bajopas.v1i1.57512](https://doi.org/10.4314/bajopas.v1i1.57512).

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

PERBAIKAN ARTIKEL SESUAI TELAAH REVIEWER

ABSTRACT

This research was conducted to determine physical quality, the nutritional quality and characteristics of antimicrobial compounds from fermented oil palm fronds (FOPF). This research was carried out using a Complete Randomized Design with 4 treatments and 4 replications. The following treatments were performed by the addition of different additives namely: 10% poultry manure, 10% cow feces, 5% urea and 5% molasses. All treatments were extracted in stages using n-hexane, ethyl acetate and methanol. The observed parameters were physical quality (pH, aroma, color, and texture), nutritional quality (**neutral detergent fiber (NDF)**, **acid detergent fiber (ADF)**, hemicellulose, cellulose and lignin), yield of extract, class of bioactive compounds and the antimicrobial activity test of extracted compounds. The data were analyzed using a one-way analysis of variance and differences between treatments were tested using Duncan's multiple range test. The addition of different additives has a very significant effect ($P < 0.01$) on the lignin content; significant effect ($P < 0.05$) on pH, aroma, color and texture; and no effect on the content of NDF, ADF, hemicellulose and cellulose. The highest yield of extract was obtained from methanol extract with the addition of poultry manure. The methanol extract on the addition of poultry manure and urea were identified as compounds in the steroid, tannin and phenolic classes. The highest antimicrobial activity test against *Escherichia coli* (12.83 mm) was obtained from FOPF extract (1%) with addition of poultry manure, while highest antimicrobial activity test against *Staphylococcus aureus* (11,67 mm) was obtained from FOPF extract (4%) with addition of molasses. **The conclusion of this research was FOPF with addition of poultry manure provides good physical quality, nutritional quality and can inhibit the growth of *E. coli* and *S. aureus* at a concentration of 1%.**

Background

The utilization of oil palm plantation waste such as oil palm fronds as feed is still limited due to the high lignin content [1, 2]. Physical-chemical-biological and combination treatments and the addition of additives on the fermentation process aim to break down lignocellulose and lignohemicellulose bonds and increase nutrient digestibility [3, 4, 5].

Palm oil fronds have antimicrobial activity [6]. Methanol extract from palm oil leaves contains tannin, alkaloids, saponins, terpenoids and flavonoids [7]; and toxicity tests on shrimp and rats showed that palm oil leaf extract is safer for consumption and recommended as a natural product [8]. This study aims to determine physical quality, the nutritional quality and characteristics of antimicrobial compounds from fermented oil palm fronds.

Methods

The research was conducted in June-November 2017, at the Laboratory of Nutrition and Chemistry at the Faculty of Agriculture and Animal Sciences of State Islamic University of

Sultan Syarif Kasim Riau Indonesia, Biotechnology Research Center of the Indonesian Sciences Institute Indonesia and Laboratory of Biota Sumatra Andalas University.

Materials

The materials used in this research were **neutral detergent soluble (NDS) solution**, **acid detergent soluble (ADS) solution**, acetone, alcohol, octanol, H_2SO_4 , methanol, ethyl acetate, hexane, filter paper, rotary vacuum evaporator, HCl, ammonia, Mayer reagent, Dragendorff reagent, ether, Liebermann Bouchard reagent, Mg powder, amyl alcohol, NaOH, $FeCl_3$, aquadest, poultry manure, cow feces, urea, molasses, plastic bags.

Research Procedure

The study was conducted through two (2) stages: namely fermentation of oil palm frond based on Febrina [9] modified with different additive (10% poultry manure, 10% cow feces, 5% urea and 5% molasses) using a Completely Randomized Design 4 treatment with 4 replications. Fermentation process for 21 days.

The data were analyzed using a one-way analysis of variance and differences between treatments were tested using Duncan's Multiple Range Test. The observed parameters were physical quality tests include pH, aroma, color and texture and nutritional quality including the content of NDF, ADF, hemi cellulose, cellulose and lignin based on Van Soest analysis using the FOSS Fibertec system. The second stage is multilevel maceration and extraction using methanol, ethyl acetate and n-hexane solvents. Phytochemical screening based on Franswort

[10] and the antibacterial activity test of oil palm fronds extract based on Handayani *et al.* [11].

Result and Discussion

The Effect of Different Additives on The Physical quality of FOPF

The addition of different additives has a significant effect ($P < 0.05$) on the physical quality of FOPF including pH, aroma, color and texture (Table 1).

Table. 1 The effect of different additives on the physical quality of FOPF

Treatment	Physical quality							
	pH		Aroma		Color		Texture	
	Value	Notes	Value	Notes	Value	Notes	Value	Notes
A	5,18 ^a ±0,10	Acid	3,00 ^b ±0,00	Acid	3,03 ^a ±0,05	yellowish green	2,00 ^a ±0,00	Rather soft
B	5,52 ^b ±0,20	Acid	4,00 ^c ±0,00	more acidic	3,00 ^a ±0,00	yellowish green	3,00 ^b ±0,00	soft
C	7,60 ^c ±0,14	Base	2,03 ^a ±0,05	Non acid	4,03 ^b ±0,05	brownish green	3,03 ^b ±0,05	soft
D	5,28 ^{ab} ±0,21	acid	5,00 ^d ±0,00	Very acid	3,00 ^a ±0,00	yellowish green	3,00 ^b ±0,00	soft

Notes : A = FOPF with the addition of poultry manure
 B = FOPF with the addition of cow feces
 C = FOPF with the addition of urea
 D = FOPF with the addition of molasses
 a,b,c means with different superscript differ ($p < 0.05$)

The lowest of pH (5.18) in treatment A is related to the availability of carbohydrates and the production of lactic acid bacteria. The same results were seen in treatment D with a pH of 5.28. Addition of poultry manure and molasses on fermentation oil palm frond aims as a source of additive which can trigger the growth of lactic acid bacteria to produce lactic acid. Increased production of lactic acid will reduce pH and inhibit the growth of pathogenic bacteria. Ozduven *et al.* [12] stated that pH is associated with lactic acid produced during the fermentation process which can inhibit the growth of pathogenic microbes. Oil palm fronds which are fermented with molasses produce pH 3.93 [3].

The process of ammonization of oil palm fronds using urea will produces the highest pH of 7.60 which is alkaline. The same results were reported by Zahari *et al.* [3], the process of

ammonization of oil palm fronds using urea resulting in a pH of 7.38. In the process of ammonization with urea will break down into ammonium ions so that the ammonization product is alkaline. Laura and Paolo [13] state that urea will break down into ammonium ions, which are alkaline and ammonization processes can break down glycoside bonds [14] so that the oil palm fronds can be degraded by rumen microbes.

In treatment A produces an aroma score of 3 with acid criteria. It has been shown that the fermentation process is going well as indicated by acid production. Aroma is related to the production of lactic acid and pH. It can be seen in Table 1 the addition of poultry manure to oil palm fronds fermentation resulting in the lowest pH (5.18) so that the products are acidic.

Treatment A, B and D produces a color that is not different ($P > 0.05$), which is natural green or

yellowish green with score of 3-3.03. Good quality silage will produce colors that are not different from the original material [15]. The occurrence of color changes from green to yellowish green indicated the anaerobic phase oxygen has run out of carbohydrates. Then it will produce CO₂, H₂O and heat. The heat produced causes a change in color from green to yellowish green.

Treatment A will produce softer textures with score 2. This shows that the addition of the poultry manure to the fermentation process can support the growth of lactic acid bacteria to produce lactic acid. Optimal production of lactic acid will reduce pH so that it inhibits the growth of pathogenic bacteria. This low pH also supports the production of lignase enzymes which function to break down lignocellulose and lignohemicellulose thereby reducing lignin

content. It is seen that treatment A produced the lowest lignin content of 19.94%. The release of lignin from cellulose and hemicellulose causes fermented products to have a softer texture.

The Effect of Different Additive on Fiber Fraction of FOPF

The addition of different additive had no effect (P>0.05) on the content of NDF, ADF, hemicellulose and cellulose but had a very significant effect (P<0.01) on the lignin content (Table 2). Rahman *et al.* [4] reported there was a decrease in the lignin content of oil palm fronds in the fermentation process using white rot fungi.

Table 2. The effect of different additive on fiber fraction of FOPF.

Treatment	Fiber Fraction (%)				
	NDF	ADF	Hemicellulose	Cellulose	Lignin
A	73,34±3,68	51,81±4,82	21,52±6,07	28,54±3,36	19,94 ^a ±1,47
B	75,39±2,03	55,03±4,23	20,35±4,46	29,57±3,09	23,15 ^b ±0,96
C	73,83±1,01	53,30±3,43	20,51±4,30	31,55±1,99	21,57 ^a ±1,58
D	74,98±3,52	54,04±4,48	20,93±6,63	31,70±4,27	22,41 ^b ±1,22

Notes: A = FOPF with the addition of poultry manure
 B = FOPF with the addition of cow feces
 C = FOPF with the addition of urea
 D = FOPF with the addition of molasses
 a,b,c means with different superscript differ (p<0.05)

Lignin is a part of plant cell walls which consists of very complex and difficult to digest bonds. The high content of lignin in feed will reduce its digestibility. The lignin content of oil palm fronds is 30.18% [2]. One of the purposes of processing feed through ammonization and fermentation in oil palm fronds is to reduce the lignin content. The addition of different additive gave a very significant effect (P <0.01) on the lignin content of FOPF. The lowest lignin content in treatment A (19.94%) was not significantly different (P> 0.05) compared to treatment C (21.57%) but was very significant (P <0.01) lower treatment B (23.15%) and treatment D (22.41%).

The lowest lignin content in treatment A (19.94%) allegedly in the oil palm frond fermentation using poultry manure will produce

higher lignase and urease enzymes than other treatments. The lignase enzyme is an extracellular enzyme that functions to degrade lignin and urease to break down urea into ammonia. Poultry manure also functions as a source of urease which will break down urea into ammonia and CO₂. The high N content in poultry manure, which is 3.63-5.30% with C/N ratio 5.8-7.6 [16], causes high urease activity so that the addition of poultry manure as a source of additive in palm frond fermentation can reduce lignin content. Thambirajah *et al.* [17] there is a positive correlation between the nitrogen content of poultry manure and the nutritional content of oil palm empty fruit bunches fermented with poultry manure. The addition of 15% poultry manure in ammoniated

rice straw can reduce fermentation time from 20 days to 5 days.

Bioactive Compounds from Extracts of FOPF

Bioactive compounds contained in extracts from FOPF can be known through maceration method with multilevel extraction. This is based on multilevel extraction can separate the bioactive compounds contained in the extract from the

FOPF in the same sample based on their level of polarity without the bioactive compounds dissolving in other solvents, therefore in this study the multilevel extraction process starts from a non-polar solvent (n-hexane) followed by a semi-polar solvent (ethyl acetate) and finally with a polar solvent (methanol).

Tabel 3. The yield of extracts FOPF

Treatment	Yield of extracts (%)		
	N heksan	Etil Asetat	Metanol
A	0,78	1,53	7,87
B	1,08	1,30	7,07
C	1,02	1,17	5,79
D	0,89	1,08	7,40

Notes : A = FOPF with the addition of poultry manure
 B = FOPF with the addition of cow feces
 C = FOPF with the addition of urea
 D = FOPF with the addition of molasses

Table 3 shows n acetate extracts FOPF produce the lowest yields ranging from 0.78-1.08% while the methanol extract produces the largest yields ranging from 5.79-7.87%. The same results were reported by Salamah *et al.* [18] the lowest yield of Taiwanese gravestone extract was from n-hexane extract (0.04%) and the highest yield was from methanol extract (0.12%).

FOPF which was extracted with methanol, produced the highest yield of 7.87% of crude compounds. This shows that the most bioactive compounds found in FOPF are polar because they can dissolve in polar solvents namely methanol. Bioactive compounds of FOPF which are non-polar and semi-polar in smaller amounts. Phytochemical analysis of bioactive compounds of extract of fermented oil palm fronds is shown in Table 4.

The extraction process uses methanol solvents in treatments A and C, produce small amounts of tannin and phenolic compounds (+). Sufficient amounts of phenolic compounds (++) were found in treatment B and D. N hexane extract in treatment A produced the lowest yield (0.78%) and methanol extract produced the highest yield (7.87%). This shows that treatment A contains polar compounds (methanol) which are greater than non-polar compounds (n hexane). Analysis of phytochemical compounds shows, treatment A contains bioactive compounds, are steroids and tannins, in small amounts (+) and phenolic compounds in sufficient quantities (++)

Table 4.

Table 4. Bioactive compounds of extract of FOPF

Class of compounds	n hexane				Etil Asetat				Metanol			
	A	B	C	D	A	B	C	D	A	B	C	D
Alkaloid	-	-	-	-	-	-	-	-	-	-	-	-
Steroid	+	+	+	+	+	+	+	+	+	+	+	+
Terpenoid	-	-	-	-	-	-	-	-	-	-	-	-
Flavanoid	-	-	-	-	-	-	-	-	-	-	-	-
Saponin	-	-	-	-	-	-	-	-	-	-	-	-
Kuinon	-	-	-	-	-	-	-	-	-	-	-	-
Tannin	-	-	-	-	-	-	-	-	+	-	+	-
Kumarin	-	-	-	-	-	-	-	-	-	-	-	-
Phenolic	-	-	-	-	-	-	-	-	+	++	+	++

Notes : A = FOPF with the addition of poultry manure
 B = FOPF with the addition of cow feces
 C = FOPF with the addition of urea
 D = FOPF with the addition of molasses
 - there are no bioactive compounds
 + in small quantities
 ++ there are in sufficient quantities

a. Steroid

Steroids are initially considered only as components of animal substances (as sex hormones, adrenal hormones, bile acids, etc.), but recently steroids have also been found in plant substances [19]. Steroids contained in methanol extract, ethyl acetate and n hexane in treatment A are thought to be adrenal hormones and sex hormones, because in poultry, metabolic waste that cannot be used again is released as excreta through cloaca. Excreta consists of feces, urine and sperm/egg cells. Bose *et al.* [20] stated that steroid compounds were detected in *Achatina fulica* (freshwater gastropods such as golden snails). Steroids are thought to increase body stamina (aphrodisiac) and anti-inflammation. Silva *et al.* [21] reported extracts from the leaves of *Agave attenuate*, containing steroid compounds that have anti-inflammatory activity.

Phytochemical analysis showed that bioactive steroid compounds were detected in all extract of FOPF (n hexane, ethyl acetate and methanol) which had different polarity. Harbone [19] states, steroids are cholesterol, which are non-polar so they can dissolve in non-polar solvents, (n-hexane), semi-polar (ethyl acetate) and polar (methanol) in small amounts (+). The steroid content in beef,

which is extracted using non-polar and polar compounds, shows no difference [22].

Bioactive compounds of steroid, which are found in extract of FOPF can be used as antimicrobials because steroids are bioactive compounds that function as anti-fungal. N hexane extract from lotus seeds contains bioactive steroid compounds and can inhibit the growth of Gram-negative bacteria (*Aeromonas hydrophila*) and Gram-positive bacteria (*Streptococcus agalactiae*) and fungi (*Saprolegnia sp.*) [23].

b. Tannin

Methanol extract from FOPF in treatments A and C contains small amounts of tannin compounds (+). The tannin compounds found in oil palm plants are reported by Sasidharan [7] in methanol extract of palm oil leaves; Febrina *et al.* [6] on the ethanol extract of oil palm fronds.

The presence of tannin compounds in methanol extract in treatments A and C shows that methanol extract from FOPF has the potential to be antimicrobial. Tannin has antibacterial properties by precipitation proteins because it is suspected that tannins have the same effect as phenolic compounds [24]. The tannin compounds found in the ethanol extracts of oil palm fronds [7] and

mangosteen peel [25] have the potential to be antimicrobial because they can inhibit the growth of *S. aureus* and *E. coli*.

c. Phenolic

Phytochemical analysis in treatment B and D showed that sufficient quantities of phenolic bioactive compounds (++) . This indicates that the extract of oil palm frond which is fermented with cow feces and molasses can be used as an antimicrobial. Phenol compounds

can be used as antibacterial because they contain aromatic rings with one or two hydroxyl groups that can change the permeability of the cytoplasmic membrane thus inhibiting bacterial growth [19,26]. Ethyl acetate extract of binahong leaves, containing polyphenols and saponins with the minimum inhibitory concentration (MIC) against *Shigella flexneri* was 8% b/v [27].

Table 5. Inhibited Zone Diameter of Extracts from FOPF

Concentration	Inhibited Zone Diameter (mm)							
	<i>E. coli</i>				<i>S. aureus</i>			
	A	B	C	D	A	B	C	D
1 %	12,83	9,00	9,00	7,67	8,00	7,67	11,33	6,30
2 %	11,67	8,67	8,83	7,33	9,16	8,67	10,16	7,67
3 %	10,50	8,66	9,00	7,50	9,67	8,00	10,16	9,16
4 %	8,50	10,33	8,00	8,00	7,83	9,33	8,00	11,67
5 %	7,33	9,00	8,83	7,50	7,83	6,67	9,00	10,00

Notes : A = extracts from FOPF with the addition of poultry manure
 B = extracts from FOPF with the addition of cow feses
 C = extracts from FOPF with the addition of urea
 D = extracts from FOPF with the addition of molasses

Antimicrobial activity test of extracts from FOPF (In Vitro)

The methanol extract from oil palm fronds which are fermented with different additive has antimicrobial activity which is characterized by the formation of inhibitory zones. The formation of inhibition zones shows that it has inhibited the growth of Gram-positive bacteria (*S. aureus*) and Gram-negative bacteria (*E. coli*), by compounds contained in methanol extract from FOPF. The inhibitory zone formed shows that bacteria have antimicrobial activity [28], thus it can be concluded that methanol extract from FOPF which are fermented with different sources of additives has antimicrobial activity that is bactericide (killing bacteria).

The inhibitory zone formed in testing the antimicrobial activity of extracts from FOPF in Gram-positive (*S.aureus*) and Gram-negative (*E. coli*) bacteria at concentrations of 1-5% as shown in Table 5.

In Table 5 shown the difference in diameter of inhibition zones in each treatment. The diameter of the inhibition zone ranges from 6.67 - 12.83 mm. The smallest diameter inhibition zone of 6.67 mm was produced at a concentration of 5% in treatment B. The largest diameter inhibition zone is 12.83 mm at a concentration of 1% in treatment A.

The ability of extracts from FOPF inhibits bacterial growth in the moderate to strong, because the diameter of the inhibition zone ranges from 6.67-12.83 mm. In accordance with the statement of Davis and Stout [29] and Mudi and Ibrahim [30] if the diameter of the inhibition zone is more than 6 mm, the extract has antimicrobial activity with a moderate to strong category.

Table 5 shows the diameter of the inhibitory zone in Gram-negative bacteria (*E. coli*) generally greater than the Gram-positive bacteria (*S. aureus*). The largest diameter of the

inhibition zone in Gram-negative bacteria (*E. coli*) namely in treatment A with a concentration of 1% was 12.83 mm. **The largest diameter of the inhibition zone in Gram-positive bacteria (*S. aureus*) namely in treatment D with a concentration of 4% is 11.67 mm.**

There was a difference in the diameter of the inhibitory zone in gram positive and Gram-negative bacteria due to differences in the structure of bacterial cell walls. The data in Table 5 shows that the antibacterial compounds of the methanol extract of FOPF were able to inhibit the growth of Gram-negative bacteria better than Gram-positive bacteria which was characterized by a larger diameter of the inhibition zone. Cell walls of Gram-positive bacteria have a simple structure compared to Gram-negative bacteria that have complex structures. The largest diameter inhibition zone (12.83 mm) for gram negative bacteria (*E. coli*) showed that the compounds contained in the methanol extract of FOPF had a greater ability to penetrate the cell walls of Gram-negative bacteria. This is because Gram-negative bacteria have a more complex structure and have more layers so that only certain compounds can penetrate it.

The minimum inhibition concentration (MIC) shows the lowest concentration of extract which can inhibit bacterial growth. The smaller the MIC value the greater the antimicrobial activity. This can be seen in treatment A at a concentration of 1% which was able to inhibit the growth of pathogenic microbes with the largest diameter of inhibition zone (12.83 mm).

Conclusion

The conclusion of this research was FOPF with addition of poultry manure provides good nutritional quality and can inhibit the growth of *E. coli* and *S. aureus* at a concentration of 1%.

Acknowledgement

Author would like to thank to the Institute for Research and Community Service of the State Islamic University of Sultan Syarif Kasim Riau which funded this study under contract number 1043/R/2017.

References

1. Febrina D, Jamarun N, Zain M, Khasrad (2016) The effects of P, S and Mg supplementation of oil palm fronds fermented by *Phanerochaete chrysosporium* on rumen fluid characteristics and microbial protein synthesis. Pakistan Journal of Nutrition 15: 299 – 304. doi: doi.org/10.3923/pjn.2016.299.304.
2. Febrina D, Jamarun N, Zain M, Khasrad (2016) Effects of calcium (Ca) and manganese (Mn) supplementation during oil palm frond fermentation by *Phanerochaete chrysosporium* on in vitro digestibility and rumen fluid characteristics. Pakistan Journal of Nutrition 15: 352 – 358. doi: doi.org/10.3923/pjn.2016.352.358.
3. Zahari WM, Hassan OA, Wong HK, Liang JB (2003) Utilization of oil palm frond - based diets for beef and dairy production in Malaysia. Asian-Australasian Journal of Animal Sciences 16 (4): 625 – 634. doi: doi.org/10.5713/ajas.2003.625.
4. Rahman MM, Lourenço M, Hassim HA, Baars JJP, Sonnenberg ASM, Cone JW, de Boever J, Fievez V (2011) Improving ruminal degradability of oil palm fronds using white rot fungi. Animal Feed Science and Technology 169 (3-4) :157 – 166. doi: doi.org/10.1016/j.anifeedsci.2011.06.014.
5. Kawamoto H, Mohamed WZ, Sukur NIM, Ali MSM, Islam Y, Oshio S (2001) Palatability, digestibility and voluntary intake of processed oil palm fronds in cattle. Japan Agricultural Research Quarterly 35; 195 – 200. doi: doi.org/10.6090/jarq.35.195.
6. Febrina D, Febriyanti R, Zam SI, Handoko J, Fatah A, Juliantoni J (2018) Antibacterial activity testing and ethanol extract characterization of oil palm fronds (*Elaeis guineensis* Jacq.). Pakistan Journal

of Nutrition 17 (9): 427 – 433. doi: 10.3923/pjn.2018.427.433.

7. Sasidharan S, Rajoo N, Rathinam X, Lachimanan YL, Rajoo A (2010) Wound healing potential of *Elaeis guineensis* Jacq. leaves in an infected albino rat model. *Molecules* 15: 3186 – 3199. doi: 10.3390/molecules15053186.

8. Syahmi ARM, Vijayarathna S, Sasidharan S, Latha LY, Kwan YP, Lau YL, Shin LN, Chen Y (2010) Acute oral toxicity and brine shrimp lethality of *Elaeis guineensis* Jacq. (oil palm leaf) methanol extract. *Molecules* 15: 8111 – 8121. doi: 10.3390/molecules15118111.

9. Febrina D (2016) Utilization of oil palm fronds biodelignification products using *Phanerochaete chrysosporium* as substitute forage on goat. PhD thesis. Andalas University, Department Animal Science.

10. Franswort NR (1996) Biological and phytochemical screenings of plant. *Journal of Pharmaceutical Science* 55 (3): 225 – 265. doi: 10.1002/jps.2600550302.

11. Handayani D, Sandrawaty N, Murniati M, Regina R (2015) Screening of endophytic bacteria isolated from marine sponge *Haliclona fascigera* for inhibition against clinical isolates of methicillin resistant *Staphylococcus aureus* (MRSA). *Journal of Applied Pharmaceutical Science* 5: 139 – 142. doi: 10.7324/JAPS.2015.50926.

12. Ozduven ML, Koc F, Polat C, Coskuntuna L (2009) The effects of lactic acid bacteria and enzyme mixture inoculants on fermentation and nutrient digestibility of sunflower silage. *Kafkas Univ Vet Fak Derg* 15 (2): 195 – 199. doi: 10.9775/kvfd.2008.91-A.

13. Laura B, Paolo G (2006) *Algae: Anatomy Biochemistry and Biotechnology*. Boca Raton, CRC Press.

14. Umiyasih U, Wina E (2008) Processing and nutritional value of corn by-product as ruminant feed. *Wartazoa* 18 (3): 127 – 136.

15. Saun RJV, Henrich AJ (Eds) (2008) *Proceedings of the Mid-Atlantic Conference: 26 May 2008; Pennsylvania*. Penn States Collage.

16. Amanullah MA, Sekar S, Muthukrishnan P (2010) Prospect and potential of poultry manure. *Asian Journal of Plant Sciences* 9 (4): 1 – 11. doi: 10.3923/ajps.2010.172.182.

17. Thambirajah JJ, Zulkali MD, Hashim MA (1995) Microbiological and biochemical changes during the composting of oil palm empty-fruit-bunches. Effect of nitrogen supplementation on the substrate. *Bioresource Technology* 52 (2): 133 – 144. doi: 10.1016/0960-8524(95)00008-3.

18. Salamah E, Ayuningrat E, Purwaningsih S (2008) Initial screening of freshwater mussel bioactive components from (*Anodonta woodiana* Lea.) from antioxidant compounds. *Buletin Teknologi Hasil Perikanan* 11 (2): 119 – 132. doi: 10.17844/jphpi.v11i2.912.

19. Harborne AJ. *Phytochemical Methods A Guide to Modern Techniques of Plant Analysis*. 3rd Edition. Dordrecht, Springer.

20. Bose R, Majumdar C, Bhattacharya S (1997) Steroids in *Achatina fulica* (bowdich): Steroid profile in haemolymph and in vitro release of steroids from endogenous precursors by ovotestis and albumen gland. *Comparative Biochemistry and Physiology* 116 (3): 179 – 182. doi: 10.1016/S0742-8413(96)00163-6.

21. Silva BP, Sousa AC, Silva GM, Mendes TP, Parente JP. A new bioactive steroidal saponin from *Agave attenuata*. *Zeitschrift für Naturforschung* 57c: 423 – 428.

22. Schmidt G, Steinhart H (2001) Impact of extraction solvents on steroid contents determined in beef. *Food Chemistry* 76 (1): 83 – 88. doi: 10.1016/S0308-8146(01)00237-0.

23. Widya DR, Suryanto D, Desrita (2014) Antimicrobial activity of lotus seeds (*Nymphaea pubescens* L.) on bacteria *Aeromonas hydrophila*, *Streptococcus agalactiae*, and *Saprolegnia* sp. *Jurnal Aquacoastmarine* 2 (1): 7 – 14.

24. Martono B, Setiyono RT (2014) Phytochemical screening of six tea genotypes. *Journal of Industrial and Beverage Crops* 1 (2): 63 – 68. doi: 10.21082/jtidp.v1n2.2014.p63-68.

25. Poelongan M, Praptiwi (2010) Antibacterial activity test of mangos teen (*Gardnia mangostana* Linn.) peel. *Media Litbang Kesehatan*. 20 (2): 65 – 69.

26. Ikalinus R, Widyastuti SK, Setiasih NLE (2015) Phytochemical screening ethanol extract skin stem moringa (*Moringa oleifera*). *Indonesia Medicus Veterinus* 4 (1): 71 – 79.

27. Wardhani LK, Sulistyani N (2012) Antibacterial activity test of ethyl acetate extract of binahong leaf (*Anredera scandens* (L.) Moq.) against *Shigella flexneri* with the profile of thin layer chromatography. *Jurnal Ilmiah Kefarmasian* 2 (1): 1 – 16.

28. Nanda. A, Saravanan M (2009) Biosynthesis of silver nanoparticles from *Staphylococcus aureus* and its antimicrobial activity against MRSA and MRSE. *Nanomedicine: Nanotechnology, Biology and Medicine* 5 (4): 452 – 456. doi: 10.1016/j.nano.2009.01.012.

29. Davis WW, Stout TR (1971) Disc plate method of microbiological antibiotic assay. I. Factors influencing variability and error. *Applied Microbiology* 22: 659 – 665.

30. Mudi SY, Ibrahim H (2008) Activity of *Bryophyllum pinnatum* S. Kurz extracts on respiratory tract pathogenic bacteria. *Bayero Journal of Pure Applied Sciences* 1 (1): 43 – 48. doi: 10.4314/bajopas.v1i1.57512.

PEMBERITAHUAN DARI EDITOR

“ACCEPT WITH REVISION”

15:54

4G+ 46%



[JTLS] Editor Decision

Inbox



Nashi Widodo 10/23/2019

to me



Mr Syukria Ikhsan Zam:

We have reached a decision regarding your submission to Journal of Tropical Life Science, "Nutritional Content and Characteristics of Antimicrobial Compounds from Fermented Oil Palm Fronds (*Elaeis guineensis* Jacq.)".

Our decision is to: accept with revision

Prof. Dr. Fahrul zaman Huyop
Universiti Teknologi Malaysia
Phone 0126849374
Fax 0126849374
fahrul@utm.my

Fahrul Huyop
Journal of Tropical Life Science
Universitas Brawijaya

PEMBERITAHUAN DARI EDITOR

“PROOF READ”

15:56

4G+ 45%



[JTLS] Galley Proof_Nutritional
Content and Characteristics of
Antimicrobial Compounds



Inbox



Nashi Widodo 1/14/2020

to me



Dear author,

Your article will be published in Journal of Tropical Life
Science Vol 10
(2020). Please:

1. Check your proof carefully and please make sure that given names and surnames of authors have been identified correctly write the full name
2. Check your corresponding author address
3. Send the Figure 2 since you mention it in the text.
4. Mark your correction using highlight and add comments, also you can send us separated list of the parts of your article that need to be corrected

Please send back your confirmation in 2 x 24 hours to this email

Thank you
Journal of Tropical Life Science
Universitas Brawijaya

Nutritional Content and Characteristics of Antimicrobial Compounds from Fermented Oil Palm Fronds (*Elaeis guineensis* Jacq)

Dewi Febrina¹, Rahmi Febriyanti², Syukria Ikhsan Zam^{2*}, Zumarni¹, Jepri Juliantoni¹,
Abdul Fatah¹

¹ Department of Animal Science, Faculty of Agriculture and Animal Sciences, State Islamic
University of Sultan Syarif Kasim Riau, Pekanbaru 28293, Indonesia

² Department of Agrotechnology, Faculty of Agriculture and Animal Sciences, State Islamic
University of Sultan Syarif Kasim Riau, Pekanbaru 28293, Indonesia

* corresponding author : email : syukria.ikhsan.zam@uin-suska.ac.id

ABSTRACT

This research was conducted to determine physical quality, the nutritional quality and characteristics of antimicrobial compounds from fermented oil palm fronds (FOPF). This research was carried out using a Complete Randomized Design with 4 treatments and 4 replications. The following treatments were performed by the addition of different additives namely: 10% poultry manure, 10% cow feces, 5% urea and 5% molasses. All treatments were extracted in stages using n-hexane, ethyl acetate and methanol. The observed parameters were physical quality (pH, aroma, color, and texture), nutritional quality (neutral detergent fiber (NDF), acid detergent fiber (ADF), hemicellulose, cellulose and lignin), yield of extract, class of bioactive compounds and the antimicrobial activity test of extracted compounds. The data were analyzed using a one-way analysis of variance and differences between treatments were tested using Duncan's multiple range test. The addition of different additives has a very significant effect ($P < 0.01$) on the lignin content; significant effect ($P < 0.05$) on pH, aroma, color and texture; and no effect on the content of NDF, ADF, hemicellulose and cellulose. The highest yield of extract was obtained from methanol extract with the addition of poultry manure. The methanol extract on the addition of poultry manure and urea were identified as compounds in the steroid, tannin and phenolic classes. The highest antimicrobial activity test against *Escherichia coli* (12.83 mm) was obtained from FOPF extract (1%) with addition of poultry manure, while highest antimicrobial activity test against *Staphylococcus aureus* (11.67 mm) was obtained from FOPF extract (4%) with addition of molasses. The conclusion of this research was FOPF with addition of poultry manure provides good physical quality, nutritional quality and can inhibit the growth of *E. coli* and *S. aureus* at a concentration of 1%.

Keywords : *Nutritional content, Antimicrobial, Oil palm fronds*

Introduction.

The utilization of oil palm plantation waste such as oil palm fronds as feed is still limited due to the high lignin content [1, 2]. Physical-chemical-biological and combination treatments and the addition of additives on the

fermentation process aim to break down lignocellulose and ligno hemicellulose bonds and increase nutrient digestibility [3, 4, 5].

Palm oil fronds have antimicrobial activity [6]. Methanol extract from palm oil leaves contains tannin, alkaloids, saponins, terpenoids

and flavonoids [7]; and toxicity tests on shrimp and rats showed that palm oil leaf extract is safe for consumption and recommended as a natural product [8]. This study aims to determine physical quality, the nutritional quality, and characteristics of anti- microbial compounds from fermented oil palm fronds.

Materials and Methods.

The research was conducted in June-November 2017, at the Laboratory of Nutrition and Chemistry at the Faculty of Agriculture and Animal Sciences of State Islamic University of Sultan Syarif Kasim Riau Indonesia, Biotechnology Research Center of the Indonesian Sciences Institute Indonesia and Laboratory of Biota Sumatra Andalas University

The study was conducted through two (2) stages: namely fermentation of oil palm frond based on Febrina [9] modified with different additive (10% poultry manure, 10% cow feces, 5% urea and 5% molasses) using a Completely Randomized Design 4 treatment with 4

replications. Fermentation process for 21 days. The data were analyzed using a one-way analysis of variance and differences between treatments were tested using Duncan's Multiple Range Test. The observed parameters were physical quality tests include pH, aroma, color and texture and nutritional quality including the content of NDF, ADF, hemi cellulose, cellulose, and lignin based on Van Soest analysis using the FOSS Fibertec system. The second stage is multilevel maceration and extraction using methanol, ethyl acetate and n-hexane solvents. Phytochemical screening based on Franswort [10] and the antibacterial activity test of oil palm fronds extract based on Handayani *et al.* [11].

Results and Discussion

The effect of different additives on the physical quality of FOPF

The addition of different additives has a significant effect ($P < 0.05$) on the physical quality of FOPF including pH, aroma, color and texture (Table 1).

Table. 1 The effect of different additives on the physical quality of FOPF

Treatment	Physical quality							
	pH		Aroma		Color		Texture	
	Value	Notes	Value	Notes	Value	Notes	Value	Notes
A	5.18 ^a ±0.10	Acid	3.00 ^b ±0.00	Acid	3.03 ^a ±0.05	yellowish green	2.00 ^a ±0.00	Rather soft
B	5.52 ^b ±0.20	Acid	4.00 ^c ±0.00	more acidic	3.00 ^a ±0.00	yellowish green	3.00 ^b ±0.00	Soft
C	7.60 ^c ±0.14	Base	2.03 ^a ±0.05	Non acid	4.03 ^b ±0.05	brownish green	3.03 ^b ±0.05	soft
D	5.28 ^{ab} ±0.21	Acid	5.00 ^d ±0.00	Very acid	3.00 ^a ±0.00	yellowish green	3.00 ^b ±0.00	Soft

Notes : A = FOPF with the addition of poultry manure
 B = FOPF with the addition of cow feces
 C = FOPF with the addition of urea
 D = FOPF with the addition of molasses
 a,b,c means with different superscript differ ($p < 0.05$)

The lowest of pH (5.18) in treatment A is related to the availability of carbohydrates and the production of lactic acid bacteria. The

same results were seen in treatment D with a pH of 5.28. Addition of poultry manure and molasses on fermentation oil palm frond aims

as a source of additive which can trigger the growth of lactic acid bacteria to produce lactic acid. Increased production of lactic acid will reduce pH and inhibit the growth of pathogenic bacteria. Ozduven *et al.* [12] stated that pH is associated with lactic acid produced during the fermentation process which can inhibit the growth of pathogenic microbes. Oil palm fronds which are fermented with molasses produce pH 3.93 [3].

The process of ammonization of oil palm fronds using urea will produce the highest pH of 7.60 which is alkaline. The same results were reported by Zahari *et al* [3], the process of ammonization of oil palm fronds using urea resulting in a pH of 7.38. In the process of ammonization with urea will break down into ammonium ions so that the ammonization product is alkaline. Laura and Paolo [13] state that urea will break down into ammonium ions, which are alkaline and ammonization processes can break down glycoside bonds [14] so that the oil palm fronds can be degraded by rumen microbes.

In treatment A produces an aroma score of 3 with acid criteria. This shows that the fermentation process is good which is characterized by acidic products were typical of fermentation. Aroma is related to the production of lactic acid and pH. It can be seen in Table 1 the addition of poultry manure to oil palm fronds fermentation resulting in the lowest pH (5.18) so that the products are acidic.

Treatment A, B and D produces a color that is not different ($P>0.05$), which is natural

green or yellowish green with score of 3-3.03. Good quality silage will produce colors that are not different from the original material [15]. The occurrence of color changes from green to yellowish green due to the anaerobic phase oxygen has run out then carbohydrates will produce CO_2 , H_2O and heat. The heat produced causes a change in color from green to yellowish green.

Treatment A will produce softer textures with score 2. This shows that the addition of the poultry manure to the fermentation process can support the growth of lactic acid bacteria to produce lactic acid. Optimal production of lactic acid will reduce pH so that it inhibits the growth of pathogenic bacteria. This low pH also supports the production of lignase enzymes which function to break down lignocellulose and lignohemicellulose thereby reducing lignin content. It is seen that treatment A produced the lowest lignin content of 19.94%. The release of lignin from cellulose and hemicellulose causes fermented products to have a softer texture.

The effect of different additive on fiber fraction of FOPF

The addition of different additive had no effect ($P>0.05$) on the content of NDF, ADF, hemicellulose and cellulose but had a very significant effect ($P<0.01$) on the lignin content. Rahman *et al* [4] reported there was a decrease in the lignin content of oil palm fronds in the fermentation process using white rot fungi.

Table 2. The effect of different additive on fiber fraction of FOPF.

Treatment	Fiber Fraction (%)				
	NDF	ADF	Hemicellulose	Cellulose	Lignin
A	73.34±3.68	51.81±4.82	21.52±6.07	28.54±3.36	19.94 ^a ±1.47
B	75.39±2.03	55.03±4.23	20.35±4.46	29.57±3.09	23.15 ^b ±0.96
C	73.83±1.01	53.30±3.43	20.51±4.30	31.55±1.99	21.57 ^a ±1.58
D	74.98±3.52	54.04±4.48	20.93±6.63	31.70±4.27	22.41 ^b ±1.22

Notes: A = FOPF with the addition of poultry manure
 B = FOPF with the addition of cow feces
 C = FOPF with the addition of urea
 D = FOPF with the addition of molasses
 a,b,c means with different superscript differ (p<0.05)

Lignin is a part of plant cell walls which consists of very complex and difficult to digest bonds. The high content of lignin in feed will reduce its digestibility. The lignin content of oil palm fronds is 30.18% [2]. One of the purposes of processing feed through ammonization and fermentation in oil palm fronds is to reduce the lignin content. The addition of different additive gave a very significant effect (P <0.01) on the lignin content of FOPF. The lowest lignin content in treatment A ie (19.94%) was not significantly different (P> 0.05) compared to treatment C (21.57%) but was very significant (P <0.01) lower treatment B (23.15%) and treatment D (22.41%).

The lowest lignin content in treatment A (19.94%) allegedly in the oil palm frond fermentation using poultry manure will produce higher lignase and urease enzymes than other treatments. The lignase enzyme is an extracellular enzyme that functions to degrade lignin and urease to break down urea into ammonia. Poultry manure also functions as a source of urease which will break down urea into ammonia and CO₂. The high N content in poultry manure, which is 3.63-5.30% with C/N ratio 5.8-7.6 [16], causes high urease activity so that the addition of poultry manure as a source of additive in palm frond fermentation can reduce lignin content.

Thambirajah *et al.* [17] there is a positive correlation between the nitrogen content of poultry manure and the nutritional content of oil palm empty fruit bunches fermented with poultry manure. The addition of 15% poultry manure in ammoniated rice straw can reduce fermentation time from 20 days to 5 days.

Bioactive compounds from extracts of FOPF
 Bioactive compounds contained in extracts from FOPF can be known through maceration method with multilevel extraction. This is based on multilevel extraction can separate the bioactive compounds contained in the extract from the FOPF in the same sample based on their level of polarity without the bioactive compounds dissolving in other solvents. Therefore in this study the multilevel extraction process starts from a non-polar solvent (n hexane) followed by a semi-polar solvent (ethyl acetate) and finally with a polar solvent (methanol).

Table 3 shows n acetate extracts FOPF produce the lowest yields ranging from 0.78-1.08% while the methanol extract produces the largest yields ranging from 5.79-7.87%. The same results were reported by Salamah et al [18] the lowest yield of Taiwanese gravestone extract was from n-hexane extract (0.04%) and the highest yield was from methanol extract (0.12%).

Tabel 3. The yield of extracts FOPF

Treatment	Yield of extracts (%)		
	N hexane	Etil Asetat	Metanol
A	0.78	1.53	7.87
B	1.08	1.30	7.07
C	1.02	1.17	5.79
D	0.89	1.08	7.40

- Notes : A = FOPF with the addition of poultry manure
 B = FOPF with the addition of cow feces
 C = FOPF with the addition of urea
 D = FOPF with the addition of molasses

FOPF which was extracted with methanol, produced the highest yield of 7.87%. This shows that the most bioactive compounds found in FOPF are polar because they can dissolve in polar solvents namely methanol.

Bioactive compounds of FOPF which are non-polar and semipolar in smaller amounts. Phytochemical analysis of bioactive compounds of extract of fermented oil palm fronds is shown in Table 4.

Table 4. Bioactive compounds of extract of FOPF

Class of compounds	n hexane				Etil Asetat				Metanol			
	A	B	C	D	A	B	C	D	A	B	C	D
Alkaloid	-	-	-	-	-	-	-	-	-	-	-	-
Steroid	+	+	+	+	+	+	+	+	+	+	+	+
Terpenoid	-	-	-	-	-	-	-	-	-	-	-	-
Flavanoid	-	-	-	-	-	-	-	-	-	-	-	-
Saponin	-	-	-	-	-	-	-	-	-	-	-	-
Kuinon	-	-	-	-	-	-	-	-	-	-	-	-
Tannin	-	-	-	-	-	-	-	-	+	-	+	-
Kumarin	-	-	-	-	-	-	-	-	-	-	-	-
Phenolic	-	-	-	-	-	-	-	-	+	++	+	++

- Notes : A = FOPF with the addition of poultry manure
 B = FOPF with the addition of cow feces
 C = FOPF with the addition of urea
 D = FOPF with the addition of molasses
 - there are no bioactive compounds
 + in small quantities
 ++ there are in sufficient quantities

The extraction process uses methanol solvents in treatments A and C, produce small amounts of tannin and phenolic compounds (+). Sufficient amounts of phenolic compounds (++) were found in treatment B and D. N hexane extract in treatment A produced the lowest yield (0.78%) and methanol extract produced the highest yield (7.87%). This shows that treatment A contains polar compounds (methanol) which are greater than non-polar compounds (n hexane). Analysis of phytochemical compounds shows, treatment A contains bioactive compounds, are steroids and

tannins, in small amounts (+) and phenolic compounds in sufficient quantities (++)

a. Steroid

Steroids are initially considered only as components of animal substances (as sex hormones, adrenal hormones, bile acids, etc.), but recently steroids have also been found in plant substances [19]. Steroids contained in methanol extract, ethyl acetate and n hexane in treatment A are thought to be adrenal hormones and sex hormones, because in poultry, metabolic waste that cannot be used again is released as excreta through cloaca.

Excreta consists of feces, urine and sperm/egg cells. Bose et al [20] stated that steroid compounds were detected in *Achatina fulica* (Freshwater gastropods such as golden snails). Steroids are thought to increase body stamina (aphrodisiac) and anti-inflammation. Silva et al. [21] reported extracts from the leaves of *Agave attenuate*, containing steroid compounds that have anti-inflammatory activity.

Phytochemical analysis showed that bioactive steroid compounds were detected in all extract of FOPF (n hexane, ethyl acetate and methanol) which had different polarity. Harbone [19] states, steroids are cholesterol, which are non-polar so they can dissolve in non-polar solvents, (n hexane), semi-polar (ethyl acetate) and polar (methanol) in small amounts (+). The steroid content in beef, which is extracted using non-polar and polar compounds, shows no difference [22].

Bioactive compounds of steroid, which are found in extract of FOPF can be used as antimicrobials because steroids are bioactive compounds that function as anti-fungal. N hexane extract from lotus seeds contains bioactive steroid compounds, and can inhibit the growth of Gram negative bacteria *Aeromonas hydrophila* and Gram positive bacteria *Streptococcus agalactiae* and *Saprolegnia* sp fungi [23].

b. Tannin

Methanol extract from FOPF in treatments A and C contains small amounts of tannin compounds (+). The tannin compounds found in oil palm plants are reported by Sasidharan [7] in methanol extract of palm oil leaves; Febrina et al [6] on the ethanol extract of oil palm fronds. The presence of tannin compounds in methanol extract in treatments A and C shows that methanol extract from FOPF has the potential to be antimicrobial. The presence of tannin compounds in methanol extract in treatments A and C shows that methanol extract from FOPF has the

potential to be antimicrobial. Tannin has antibacterial properties by precipitation proteins because it is suspected that tannins have the same effect as phenolic compounds [24]. The tannin compounds found in the ethanol extracts of oil palm fronds [7] and mangosteen peel [25] have the potential to be antimicrobial because they can inhibit the growth of *S. aureus* and *E. coli*.

c. Fenolic

Phytochemical analysis in treatment B and D showed that sufficient quantities of phenolic bioactive compounds (++). This indicates that the extract of oil palm frond which is fermented with cow feces and molasses can be used as an antimicrobial. Phenol compounds can be used as antibacterial because they contain aromatic rings with one or two hydroxyl groups that can change the permeability of the cytoplasmic membrane thus inhibiting bacterial growth [16, 22]. Ethyl acetate extract of *Binahong* leaves, containing polyphenols and saponins with the Minimum Inhibitory Concentration (MIC) against *Shigella flexneri* was 8% b/v [27].

Antimicrobial activity test of extracts from FOPF (In Vitro)

The methanol extract from oil palm fronds which are fermented with different additive has antimicrobial activity which is characterized by the formation of inhibitory zones. The formation of inhibition zones shows that it has inhibited the growth of gram-positive bacteria (*S. aureus*) and gram-negative bacteria (*E. coli*), by compounds contained in methanol extract from FOPF. The inhibitory zone formed shows that bacteria have antimicrobial activity [28], thus it can be concluded that methanol extract from FOPF which are fermented with different sources of additives has antimicrobial activity that is bacteriocide (killing bacteria). The inhibitory zone formed in testing the antimicrobial activity of extracts from FOPF in Gram

positive (*S.aureus*) and Gram negative (*E.coli*) bacteria at concentrations of 1-5% as shown in

Table 5.

Tabel 5. Inhibited Zone Diameter of Extracts from FOPF

Concentration	Inhibited Zone Diameter (mm)							
	<i>E. coli</i>				<i>S. aureus</i>			
	A	B	C	D	A	B	C	D
1 %	12.83	9.00	9.00	7.67	8.00	7.67	11.33	6.30
2 %	11.67	8.67	8.83	7.33	9.16	8.67	10.16	7.67
3 %	10.50	8.66	9.00	7.50	9.67	8.00	10.16	9.16
4 %	8.50	10.33	8.00	8.00	7.83	9.33	8.00	11.67
5 %	7.33	9.00	8.83	7.50	7.83	6.67	9.00	10.00

Notes : A = extracts from FOPF with the addition of poultry manure
 B = extracts from FOPF with the addition of cow feses
 C = extracts from FOPF with the addition of urea
 D = extracts from FOPF with the addition of molasses

Table 5 shows the difference in diameter of inhibition zones in each treatment. The diameter of the inhibition zone ranges from 6.67 - 12.83 mm. The smallest diameter inhibition zone of 6.67 mm was produced at a concentration of 5% in treatment B. The largest diameter inhibition zone is 12.83 mm at a concentration of 1% in treatment A.

The ability of extracts from FOPF inhibits bacterial growth in the moderate to strong because the diameter of the inhibition zone ranges from 6.67-12.83 mm. In accordance with the statement of Davis and Stout [29] and Mudi and Ibrahim [30] if the diameter of the inhibition zone is more than 6 mm, the extract has antimicrobial activity with a moderate to strong category.

Table 5 shows the diameter of the inhibitory zone in gram negative bacteria (*Escherichia coli*) generally greater than the gram-positive bacteria (*S. aureus*). The largest diameter of the inhibition zone in gram negative bacteria (*E. coli*) namely in treatment A with a concentration of 1% was 12.83 mm. The largest diameter of the inhibition zone in gram-positive bacteria (*S. aureus*) namely in treatment D with a concentration of 4% is 11.67 mm.

There was a difference in the diameter of the inhibitory zone in gram positive and gram negative bacteria due to differences in the structure of bacterial cell walls. The data in Table 5 shows that the antibacterial compounds of the methanol extract of FOPF were able to inhibit the growth of gram negative bacteria better than gram-positive bacteria which was characterized by a larger diameter of the inhibition zone. Cell walls of gram-positive bacteria have a simple structure compared to gram negative bacteria that have complex structures. The largest diameter inhibition zone (12.83 mm) for gram negative bacteria (*E. coli*) showed that the compounds contained in the methanol extract of FOPF had a greater ability to penetrate the cell walls of gram negative bacteria. This is because gram negative bacteria have a more complex structure and have more layers so that only certain compounds can penetrate it.

The Minimum Inhibition Concentration (MIC) shows the lowest concentration of extract which can inhibit bacterial growth. The smaller the MIC value the greater the antimicrobial activity. This can be seen in treatment A at a concentration of 1% which was able to inhibit the growth of pathogenic

microbes with the largest diameter of inhibition zone (12.83 mm).

Conclusion.

The conclusion of this research was FOPF with addition of poultry manure provides good nutritional quality and can inhibit the growth of *E. coli* and *S. aureus* at a concentration of 1%.

Acknowledgment.

Author would like to thank to the Institute for Research and Community Service of the State Islamic University of Sultan Syarif Kasim Riau which funded this study under contract number 1043/R/2017.

References

1. Febrina D, Jamarun N, Zain M, Khasrad (2016) The effects of P, S and Mg supplementation of oil palm fronds fermented by *Phanerochaete chrysosporium* on rumen fluid characteristics and microbial protein synthesis. Pakistan Journal of Nutrition 15: 299 – 304. doi: 10.3923/pjn.2016.299.304
2. Febrina D, Jamarun N, Zain M, Khasrad. (2016) Effects of calcium (Ca) and manganese (Mn) supplementation during oil palm frond fermentation by *Phanerochaete chrysosporium* on in vitro digestibility and rumen fluid characteristics. Pakistan Journal of Nutrition 15: 352-358. doi: 10.3923/pjn.2016.352.358.
3. Zahari WM, Hassan OA, Wong HK, Liang JB (2003) Utilization of oil palm frond - based diets for beef and dairy production in Malaysia. Asian-Australasian Journal of Animal Sciences 16 (4): 625 – 634. doi: 10.5713/ajas.2003.625.
4. Rahman MM, Lourenço M, Hassim HA, Baars JJP, Sonnenberg ASM, Cone JW, de Boever J, Fievez V (2011) Improving ruminal degradability of oil palm fronds using white rot fungi. *Animal Feed Science and Technology* 169 (3-4) :157 – 166. doi: 10.1016/j.anifeedsci.2011.06.014.
5. Kawamoto H, Mohamed WZ, Sukur NIM, Ali MSM, Islam Y, Oshio S (2001) Palatability, digestibility and voluntary intake of processed oil palm fronds in cattle. Japan Agricultural Research Quarterly 35; 195 – 200. doi: 10.6090/jarq.35.195.
6. Febrina D, Febriyanti R, Zam SI, Handoko J, Fatah A, Juliantoni J (2018) Antibacterial activity testing and ethanol extract characterization of oil palm fronds (*Elaeis guineensis* Jacq.). Pakistan Journal of Nutrition 17 (9): 427 – 433. doi: 10.3923/pjn.2018.427.433.
7. Sasidharan S, Rajoo N, Rathinam X, Lachimanan YL, Rajoo A (2010) Wound healing potential of *Elaeis guineensis* Jacq. leaves in an infected albino rat model. Molecules 15: 3186 – 3199. doi: 10.3390/molecules15053186.
8. Syahmi, A.R.M.; Vijayarathna, S.; Sasidharan, S.; Latha, L.Y.; Kwan, Y.P.; Lau, Y.L.; Shin, L.N.; Chen, Y (2010). Acute oral toxicity and brine shrimp lethality of *Elaeis guineensis* jacq., (oil palm leaf) methanol extract. Molecules; 15: 8111–8121. doi:10.3390/molecules1511811.
9. Febrina, D (2016) Utilization of oil palm fronds biodelignification products using *Phanerochaete chrysosporium* as substitute forage on goat. Ph.D. Thesis, Faculty of Animal Science, University of Andalas, Padang, Indonesia.

10. Franswort, N. R (1996) Biological and Phytochemical Screenings of Plant. *J. Pharm. Sci.*, 1996; 55 (3): 225-265. doi: 10.1002/jps.2600550302.
11. Handayani, D., N. Sandrawaty, M. Murniati and R. Regina (2015) Screening of endophytic bacteria isolated from marine sponge *Haliclona fascigera* for inhibition against clinical isolates of methicillin resistant *Staphylococcus aureus* (MRSA). *J. Applied Pharmaceut. Sci* 5: 139-142. doi: 10.7324/JAPS.2015.50926.
12. Ozduven. M. L., F. KOC., C. Polat., L. Coskuntuna (2009) The Effects of Lactic Acid Bacteria and Enzyme Mixture Inoculants on Fermentation and Nutrient Digestibility of Sunflower Silage. *Kafkas Univ Vet Fak Derg* 15 (2): 195-199. doi:10.9775/kvfd.2008.91-A.
13. Laura. B and G. Paolo (2006) *Algae. Anatomy Biochemistry and Biotechnology*. Boca Raton. CRC Press.
14. Umiyasih. U dan E. Wina. Processing and Nutritional Value of Corn by-Product as Ruminant Feed. *Wartazoa*. 2008; 18(3):127-136.
15. Saun RJV, Henrich A.J (Eds) (2008) *Proceedings of the Mid-Atlantic Conference: Pennsylvania, 26 May 2008*. Penn State's Collage.
16. Amanullah MA, Sekar S, Muthukrishnan P (2010) Prospect and Potential of Poultry Manure. *Asian Journal of Plant Sciences*. 9(4): 1 – 11. doi: 10.3923/ajps.2010.172.182.
17. Thambirajah JJ, Zulkali MD, Hashim MA (1995) Microbiological and biochemical changes during the composting of oil palm empty-fruit-bunches. Effect of nitrogen supplementation on the substrate. *Bioresource Technology*. 1995; 52(2): 133-144. doi: [10.1016/0960-8524\(95\)00008-3](https://doi.org/10.1016/0960-8524(95)00008-3).
18. Salamah E, Ayuningrat E, Purwaningsih S (2008) Initial Screening of Fresh Water Mussel Bioactive Components from (*Anodonta woodiana* Lea.) from Antioxidant Compounds. *Buletin Teknologi Hasil Perikanan*.11(2):119-132. doi: [10.17844/jphpi.v11i2.912](https://doi.org/10.17844/jphpi.v11i2.912).
19. Harborne AJ (1998) *Phytochemical Methods A Guide to Modern Techniques of Plant Analysis*. 3rd Edition. Dordrecht, Springer.
20. Bose R, Majumdar C, Bhattacharya S (1997) Steroids in *Achatina fulica* (Bowdich): steroid profile in haemolymph and in vitro release of steroids from endogenous precursors by ovotestis and albumen gland. *Comparative Biochemistry and Physiology* 116 (3): 179 – 182. doi: [10.1016/S0742-8413\(96\)00163-6](https://doi.org/10.1016/S0742-8413(96)00163-6).
21. Silva BP, Sousa AC, Silva GM, Mendes TP, Parente JP. A new bioactive steroidal saponin from *Agave attenuata*. *Z Naturforsch* 2002; 57c:423-428.
22. Schmidt G, Steinhart H (2001) Impact of extraction solvents on steroid contents determined in beef. *Journal of Food Chemistry*. 76:83-88. doi:10.1016/S0308-8146(01)00237-0.
23. Widya DR, Suryanto D, Desrita (2014) Antimicrobial Activity of Lotus Seeds (*Nymphaea pubescens* L.) on Bacteria *Aeromonas hydrophila*, *Streptococcus agalactiae*, and *Saprolegnia* sp. *Jurnal Aquacoastmarine* 2(1).7-14.

24. Martono B, Setiyono RT (2014) Phytochemical screening of six tea Genotypes Journal of Industrial and Beverage Crops 1 (2): 63 – 68. doi: [10.21082/jtidp.v1n2.2014.p63-68](https://doi.org/10.21082/jtidp.v1n2.2014.p63-68).
25. Poelongan M, Praptiwi (2010). Antibacterial activity test of mangos teen (*Gardnia mangostana Linn*) Peel. Media Litbang Kesehatan 20(2):65-69.
26. Ikalinus R, Widyastuti SK, Setiasih NLE (2015) Phytochemical screening ethanol extract skin stem moringa (*Moringa oleifera*). Indonesia Medicus Veterinus 4(1):71-79.
27. Wardhani LK, Sulistyani N (2012) Antibacterial activity test of ethyl acetate extract of binahong leaf (*Anredera scandens (L.) Moq.*) against *Shigella flexneri* with the profile of thin layer chromatography. Jurnal Ilmiah Kefarmasian 2(1):1-16.
28. Nanda A, Saravanan M (2009) Biosynthesis of silver nanoparticles. from *Staphylococcus aureus* and its antimicrobial activity against MRSA and MRSE. Nanomedicine: Nanotechnology, Biology and Medicine 5(4): 452-456. doi:10.1016.j.nano.2009.01.012.
29. Davis WW, Stout TR (1971), Disc plate method of microbiological antibiotic assay. I. Factors influencing variability and error. Applied Microbiol 22: 659-665.
30. Mudi SY, Ibrahim H (2008) Activity of *Bryophyllum pinnatum S. Kurz* extracts on respiratory tract pathogenic bacteria. Bayero Journal of Pure Applied Sciences 1: 43-48. doi:10.4314/bajopas.v1i1.57512.

ARTIKEL YANG SUDAH DICEK OLEH PENULIS

Nutritional Content and Characteristics of Antimicrobial Compounds from Fermented Oil Palm Fronds (*Elaeis guineensis* Jacq)

Dewi Febrina¹, Rahmi Febriyanti², Syukria Ikhsan Zam^{2*}, Zumarni¹, Jepri Juliantoni¹,
Abdul Fatah¹

¹ Department of Animal Science, Faculty of Agriculture and Animal Sciences, State Islamic
University of Sultan Syarif Kasim Riau, Pekanbaru 28293, Indonesia

² Department of Agrotechnology, Faculty of Agriculture and Animal Sciences, State Islamic
University of Sultan Syarif Kasim Riau, Pekanbaru 28293, Indonesia

* corresponding author : email : syukria.ikhsan.zam@uin-suska.ac.id

ABSTRACT

This research was conducted to determine physical quality, the nutritional quality and characteristics of antimicrobial compounds from fermented oil palm fronds (FOPF). This research was carried out using a Complete Randomized Design with 4 treatments and 4 replications. The following treatments were performed by the addition of different additives namely: 10% poultry manure, 10% cow feces, 5% urea and 5% molasses. All treatments were extracted in stages using n-hexane, ethyl acetate and methanol. The observed parameters were physical quality (pH, aroma, color, and texture), nutritional quality (neutral detergent fiber (NDF), acid detergent fiber (ADF), hemicellulose, cellulose and lignin), yield of extract, class of bioactive compounds and the antimicrobial activity test of extracted compounds. The data were analyzed using a one-way analysis of variance and differences between treatments were tested using Duncan's multiple range test. The addition of different additives has a very significant effect ($P < 0.01$) on the lignin content; significant effect ($P < 0.05$) on pH, aroma, color and texture; and no effect on the content of NDF, ADF, hemicellulose and cellulose. The highest yield of extract was obtained from methanol extract with the addition of poultry manure. The methanol extract on the addition of poultry manure and urea were identified as compounds in the steroid, tannin and phenolic classes. The highest antimicrobial activity test against *Escherichia coli* (12.83 mm) was obtained from FOPF extract (1%) with addition of poultry manure, while highest antimicrobial activity test against *Staphylococcus aureus* (11.67 mm) was obtained from FOPF extract (4%) with addition of molasses. The conclusion of this research was FOPF with addition of poultry manure provides good physical quality, nutritional quality and can inhibit the growth of *E. coli* and *S. aureus* at a concentration of 1%.

Keywords : *Nutritional content, Antimicrobial, Oil palm fronds*

Introduction.

The utilization of oil palm plantation waste such as oil palm fronds as feed is still limited due to the high lignin content [1, 2]. Physical-chemical-biological and combination treatments and the addition of additives on the

fermentation process aim to break down lignocellulose and ligno hemicellulose bonds and increase nutrient digestibility [3, 4, 5].

Palm oil fronds have antimicrobial activity [6]. Methanol extract from palm oil leaves contains tannin, alkaloids, saponins, terpenoids

and flavonoids [7]; and toxicity tests on shrimp and rats showed that palm oil leaf extract is safe for consumption and recommended as a natural product [8]. This study aims to determine physical quality, the nutritional quality, and characteristics of anti- microbial compounds from fermented oil palm fronds.

Materials and Methods.

The research was conducted in June-November 2017, at the Laboratory of Nutrition and Chemistry at the Faculty of Agriculture and Animal Sciences of State Islamic University of Sultan Syarif Kasim Riau Indonesia, Biotechnology Research Center of the Indonesian Sciences Institute Indonesia and Laboratory of Biota Sumatra Andalas University

The study was conducted through two (2) stages: namely fermentation of oil palm frond based on Febrina [9] modified with different additive (10% poultry manure, 10% cow feces, 5% urea and 5% molasses) using a Completely Randomized Design 4 treatment with 4

replications. Fermentation process for 21 days. The data were analyzed using a one-way analysis of variance and differences between treatments were tested using Duncan's Multiple Range Test. The observed parameters were physical quality tests include pH, aroma, color and texture and nutritional quality including the content of NDF, ADF, hemi cellulose, cellulose, and lignin based on Van Soest analysis using the FOSS Fibertec system. The second stage is multilevel maceration and extraction using methanol, ethyl acetate and n-hexane solvents. Phytochemical screening based on Franswort [10] and the antibacterial activity test of oil palm fronds extract based on Handayani *et al.* [11].

Results and Discussion

The effect of different additives on the physical quality of FOPF

The addition of different additives has a significant effect ($P < 0.05$) on the physical quality of FOPF including pH, aroma, color and texture (Table 1).

Table. 1 The effect of different additives on the physical quality of FOPF

Treatment	Physical quality							
	pH		Aroma		Color		Texture	
	Value	Notes	Value	Notes	Value	Notes	Value	Notes
A	5.18 ^a ±0.10	Acid	3.00 ^b ±0.00	Acid	3.03 ^a ±0.05	yellowish green	2.00 ^a ±0.00	Rather soft
B	5.52 ^b ±0.20	Acid	4.00 ^c ±0.00	more acidic	3.00 ^a ±0.00	yellowish green	3.00 ^b ±0.00	Soft
C	7.60 ^c ±0.14	Base	2.03 ^a ±0.05	Non acid	4.03 ^b ±0.05	brownish green	3.03 ^b ±0.05	soft
D	5.28 ^{ab} ±0.21	Acid	5.00 ^d ±0.00	Very acid	3.00 ^a ±0.00	yellowish green	3.00 ^b ±0.00	Soft

Notes : A = FOPF with the addition of poultry manure
 B = FOPF with the addition of cow feces
 C = FOPF with the addition of urea
 D = FOPF with the addition of molasses
 a,b,c means with different superscript differ ($p < 0.05$)

The lowest of pH (5.18) in treatment A is related to the availability of carbohydrates and the production of lactic acid bacteria. The

same results were seen in treatment D with a pH of 5.28. Addition of poultry manure and molasses on fermentation oil palm frond aims

as a source of additive which can trigger the growth of lactic acid bacteria to produce lactic acid. Increased production of lactic acid will reduce pH and inhibit the growth of pathogenic bacteria. Ozduven *et al.* [12] stated that pH is associated with lactic acid produced during the fermentation process which can inhibit the growth of pathogenic microbes. Oil palm fronds which are fermented with molasses produce pH 3.93 [3].

The process of ammonization of oil palm fronds using urea will produce the highest pH of 7.60 which is alkaline. The same results were reported by Zahari *et al* [3], the process of ammonization of oil palm fronds using urea resulting in a pH of 7.38. In the process of ammonization with urea will break down into ammonium ions so that the ammonization product is alkaline. Laura and Paolo [13] state that urea will break down into ammonium ions, which are alkaline and ammonization processes can break down glycoside bonds [14] so that the oil palm fronds can be degraded by rumen microbes.

In treatment A produces an aroma score of 3 with acid criteria. This shows that the fermentation process is good which is characterized by acidic products were typical of fermentation. Aroma is related to the production of lactic acid and pH. It can be seen in Table 1 the addition of poultry manure to oil palm fronds fermentation resulting in the lowest pH (5.18) so that the products are acidic.

Treatment A, B and D produces a color that is not different ($P>0.05$), which is natural

green or yellowish green with score of 3-3.03. Good quality silage will produce colors that are not different from the original material [15]. The occurrence of color changes from green to yellowish green due to the anaerobic phase oxygen has run out then carbohydrates will produce CO_2 , H_2O and heat. The heat produced causes a change in color from green to yellowish green.

Treatment A will produce softer textures with score 2. This shows that the addition of the poultry manure to the fermentation process can support the growth of lactic acid bacteria to produce lactic acid. Optimal production of lactic acid will reduce pH so that it inhibits the growth of pathogenic bacteria. This low pH also supports the production of lignase enzymes which function to break down lignocellulose and lignohemicellulose thereby reducing lignin content. It is seen that treatment A produced the lowest lignin content of 19.94%. The release of lignin from cellulose and hemicellulose causes fermented products to have a softer texture.

The effect of different additive on fiber fraction of FOPF

The addition of different additive had no effect ($P>0.05$) on the content of NDF, ADF, hemicellulose and cellulose but had a very significant effect ($P<0.01$) on the lignin content. Rahman *et al* [4] reported there was a decrease in the lignin content of oil palm fronds in the fermentation process using white rot fungi.

Table 2. The effect of different additive on fiber fraction of FOPF.

Treatment	Fiber Fraction (%)				
	NDF	ADF	Hemicellulose	Cellulose	Lignin
A	73.34±3.68	51.81±4.82	21.52±6.07	28.54±3.36	19.94 ^a ±1.47
B	75.39±2.03	55.03±4.23	20.35±4.46	29.57±3.09	23.15 ^b ±0.96
C	73.83±1.01	53.30±3.43	20.51±4.30	31.55±1.99	21.57 ^a ±1.58
D	74.98±3.52	54.04±4.48	20.93±6.63	31.70±4.27	22.41 ^b ±1.22

Notes: A = FOPF with the addition of poultry manure
 B = FOPF with the addition of cow feces
 C = FOPF with the addition of urea
 D = FOPF with the addition of molasses
 a,b,c means with different superscript differ (p<0.05)

Lignin is a part of plant cell walls which consists of very complex and difficult to digest bonds. The high content of lignin in feed will reduce its digestibility. The lignin content of oil palm fronds is 30.18% [2]. One of the purposes of processing feed through ammonization and fermentation in oil palm fronds is to reduce the lignin content. The addition of different additive gave a very significant effect (P <0.01) on the lignin content of FOPF. The lowest lignin content in treatment A ie (19.94%) was not significantly different (P> 0.05) compared to treatment C (21.57%) but was very significant (P <0.01) lower treatment B (23.15%) and treatment D (22.41%).

The lowest lignin content in treatment A (19.94%) allegedly in the oil palm frond fermentation using poultry manure will produce higher lignase and urease enzymes than other treatments. The lignase enzyme is an extracellular enzyme that functions to degrade lignin and urease to break down urea into ammonia. Poultry manure also functions as a source of urease which will break down urea into ammonia and CO₂. The high N content in poultry manure, which is 3.63-5.30% with C/N ratio 5.8-7.6 [16], causes high urease activity so that the addition of poultry manure as a source of additive in palm frond fermentation can reduce lignin content.

Thambirajah *et al.* [17] there is a positive correlation between the nitrogen content of poultry manure and the nutritional content of oil palm empty fruit bunches fermented with poultry manure. The addition of 15% poultry manure in ammoniated rice straw can reduce fermentation time from 20 days to 5 days.

Bioactive compounds from extracts of FOPF

Bioactive compounds contained in extracts from FOPF can be known through maceration method with multilevel extraction. This is based on multilevel extraction can separate the bioactive compounds contained in the extract from the FOPF in the same sample based on their level of polarity without the bioactive compounds dissolving in other solvents. Therefore in this study the multilevel extraction process starts from a non-polar solvent (n hexane) followed by a semi-polar solvent (ethyl acetate) and finally with a polar solvent (methanol).

Table 3 shows n acetate extracts FOPF produce the lowest yields ranging from 0.78-1.08% while the methanol extract produces the largest yields ranging from 5.79-7.87%. The same results were reported by Salamah et al [18] the lowest yield of Taiwanese gravestone extract was from n-hexane extract (0.04%) and the highest yield was from methanol extract (0.12%).

Tabel 3. The yield of extracts FOPF

Treatment	Yield of extracts (%)		
	N hexane	Etil Asetat	Metanol
A	0.78	1.53	7.87
B	1.08	1.30	7.07
C	1.02	1.17	5.79
D	0.89	1.08	7.40

- Notes : A = FOPF with the addition of poultry manure
 B = FOPF with the addition of cow feces
 C = FOPF with the addition of urea
 D = FOPF with the addition of molasses

FOPF which was extracted with methanol, produced the highest yield of 7.87%. This shows that the most bioactive compounds found in FOPF are polar because they can dissolve in polar solvents namely methanol.

Bioactive compounds of FOPF which are non-polar and semipolar in smaller amounts. Phytochemical analysis of bioactive compounds of extract of fermented oil palm fronds is shown in Table 4.

Table 4. Bioactive compounds of extract of FOPF

Class of compounds	n hexane				Etil Asetat				Metanol			
	A	B	C	D	A	B	C	D	A	B	C	D
Alkaloid	-	-	-	-	-	-	-	-	-	-	-	-
Steroid	+	+	+	+	+	+	+	+	+	+	+	+
Terpenoid	-	-	-	-	-	-	-	-	-	-	-	-
Flavanoid	-	-	-	-	-	-	-	-	-	-	-	-
Saponin	-	-	-	-	-	-	-	-	-	-	-	-
Kuinon	-	-	-	-	-	-	-	-	-	-	-	-
Tannin	-	-	-	-	-	-	-	-	+	-	+	-
Kumarin	-	-	-	-	-	-	-	-	-	-	-	-
Phenolic	-	-	-	-	-	-	-	-	+	++	+	++

- Notes : A = FOPF with the addition of poultry manure
 B = FOPF with the addition of cow feces
 C = FOPF with the addition of urea
 D = FOPF with the addition of molasses
 - there are no bioactive compounds
 + in small quantities
 ++ there are in sufficient quantities

The extraction process uses methanol solvents in treatments A and C, produce small amounts of tannin and phenolic compounds (+). Sufficient amounts of phenolic compounds (++) were found in treatment B and D. N hexane extract in treatment A produced the lowest yield (0.78%) and methanol extract produced the highest yield (7.87%). This shows that treatment A contains polar compounds (methanol) which are greater than non-polar compounds (n hexane). Analysis of phytochemical compounds shows, treatment A contains bioactive compounds, are steroids and

tannins, in small amounts (+) and phenolic compounds in sufficient quantities (++)

a. Steroid

Steroids are initially considered only as components of animal substances (as sex hormones, adrenal hormones, bile acids, etc.), but recently steroids have also been found in plant substances [19]. Steroids contained in methanol extract, ethyl acetate and n hexane in treatment A are thought to be adrenal hormones and sex hormones, because in poultry, metabolic waste that cannot be used again is released as excreta through cloaca.

Excreta consists of feces, urine and sperm/egg cells. Bose et al [20] stated that steroid compounds were detected in *Achatina fulica* (Freshwater gastropods such as golden snails). Steroids are thought to increase body stamina (aphrodisiac) and anti-inflammation. Silva et al. [21] reported extracts from the leaves of *Agave attenuate*, containing steroid compounds that have anti-inflammatory activity.

Phytochemical analysis showed that bioactive steroid compounds were detected in all extract of FOPF (n hexane, ethyl acetate and methanol) which had different polarity. Harbone [19] states, steroids are cholesterol, which are non-polar so they can dissolve in non-polar solvents, (n hexane), semi-polar (ethyl acetate) and polar (methanol) in small amounts (+). The steroid content in beef, which is extracted using non-polar and polar compounds, shows no difference [22].

Bioactive compounds of steroid, which are found in extract of FOPF can be used as antimicrobials because steroids are bioactive compounds that function as anti-fungal. N hexane extract from lotus seeds contains bioactive steroid compounds, and can inhibit the growth of Gram negative bacteria *Aeromonas hydrophila* and Gram positive bacteria *Streptococcus agalactiae* and *Saprolegnia* sp fungi [23].

b. Tannin

Methanol extract from FOPF in treatments A and C contains small amounts of tannin compounds (+). The tannin compounds found in oil palm plants are reported by Sasidharan [7] in methanol extract of palm oil leaves; Febrina et al [6] on the ethanol extract of oil palm fronds. The presence of tannin compounds in methanol extract in treatments A and C shows that methanol extract from FOPF has the potential to be antimicrobial. The presence of tannin compounds in methanol extract in treatments A and C shows that methanol extract from FOPF has the

potential to be antimicrobial. Tannin has antibacterial properties by precipitation proteins because it is suspected that tannins have the same effect as phenolic compounds [24]. The tannin compounds found in the ethanol extracts of oil palm fronds [7] and mangosteen peel [25] have the potential to be antimicrobial because they can inhibit the growth of *S. aureus* and *E. coli*.

c. Fenolic

Phytochemical analysis in treatment B and D showed that sufficient quantities of phenolic bioactive compounds (++). This indicates that the extract of oil palm frond which is fermented with cow feces and molasses can be used as an antimicrobial. Phenol compounds can be used as antibacterial because they contain aromatic rings with one or two hydroxyl groups that can change the permeability of the cytoplasmic membrane thus inhibiting bacterial growth [16, 22]. Ethyl acetate extract of Binahong leaves, containing polyphenols and saponins with the Minimum Inhibitory Concentration (MIC) against *Shigella flexneri* was 8% b/v [27].

Antimicrobial activity test of extracts from FOPF (In Vitro)

The methanol extract from oil palm fronds which are fermented with different additive has antimicrobial activity which is characterized by the formation of inhibitory zones. The formation of inhibition zones shows that it has inhibited the growth of gram-positive bacteria (*S. aureus*) and gram-negative bacteria (*E. coli*), by compounds contained in methanol extract from FOPF. The inhibitory zone formed shows that bacteria have antimicrobial activity [28], thus it can be concluded that methanol extract from FOPF which are fermented with different sources of additives has antimicrobial activity that is bacteriocide (killing bacteria). The inhibitory zone formed in testing the antimicrobial activity of extracts from FOPF in Gram

positive (*S.aureus*) and Gram negative (*E.coli*) bacteria at concentrations of 1-5% as shown in

Table 5.

Table 5. Inhibited Zone Diameter of Extracts from FOPF

Concentration	Inhibited Zone Diameter (mm)							
	<i>E. coli</i>				<i>S. aureus</i>			
	A	B	C	D	A	B	C	D
1 %	12.83	9.00	9.00	7.67	8.00	7.67	11.33	6.30
2 %	11.67	8.67	8.83	7.33	9.16	8.67	10.16	7.67
3 %	10.50	8.66	9.00	7.50	9.67	8.00	10.16	9.16
4 %	8.50	10.33	8.00	8.00	7.83	9.33	8.00	11.67
5 %	7.33	9.00	8.83	7.50	7.83	6.67	9.00	10.00

Notes : A = extracts from FOPF with the addition of poultry manure
 B = extracts from FOPF with the addition of cow feses
 C = extracts from FOPF with the addition of urea
 D = extracts from FOPF with the addition of molasses

Table 5 shows the difference in diameter of inhibition zones in each treatment. The diameter of the inhibition zone ranges from 6.67 - 12.83 mm. The smallest diameter inhibition zone of 6.67 mm was produced at a concentration of 5% in treatment B. The largest diameter inhibition zone is 12.83 mm at a concentration of 1% in treatment A.

The ability of extracts from FOPF inhibits bacterial growth in the moderate to strong because the diameter of the inhibition zone ranges from 6.67-12.83 mm. In accordance with the statement of Davis and Stout [29] and Mudi and Ibrahim [30] if the diameter of the inhibition zone is more than 6 mm, the extract has antimicrobial activity with a moderate to strong category.

Table 5 shows the diameter of the inhibitory zone in gram negative bacteria (*Escherichia coli*) generally greater than the gram-positive bacteria (*S. aureus*). The largest diameter of the inhibition zone in gram negative bacteria (*E. coli*) namely in treatment A with a concentration of 1% was 12.83 mm. The largest diameter of the inhibition zone in gram-positive bacteria (*S. aureus*) namely in

treatment D with a concentration of 4% is 11.67 mm.

There was a difference in the diameter of the inhibitory zone in gram positive and gram negative bacteria due to differences in the structure of bacterial cell walls. The data in Table 5 shows that the antibacterial compounds of the methanol extract of FOPF were able to inhibit the growth of gram negative bacteria better than gram-positive bacteria which was characterized by a larger diameter of the inhibition zone. Cell walls of gram-positive bacteria have a simple structure compared to gram negative bacteria that have complex structures. The largest diameter inhibition zone (12.83 mm) for gram negative bacteria (*E. coli*) showed that the compounds contained in the methanol extract of FOPF had a greater ability to penetrate the cell walls of gram negative bacteria. This is because gram negative bacteria have a more complex structure and have more layers so that only certain compounds can penetrate it.

The Minimum Inhibition Concentration (MIC) shows the lowest concentration of extract which can inhibit bacterial growth. The smaller the MIC value the greater the

antimicrobial activity. This can be seen in treatment A at a concentration of 1% which was able to inhibit the growth of pathogenic microbes with the largest diameter of inhibition zone (12.83 mm).

Conclusion.

The conclusion of this research was FOPF with addition of poultry manure provides good nutritional quality and can inhibit the growth of *E. coli* and *S. aureus* at a concentration of 1%.

Acknowledgment.

Author would like to thank to the Institute for Research and Community Service of the State Islamic University of Sultan Syarif Kasim Riau which funded this study under contract number 1043/R/2017.

References

1. Febrina D, Jamarun N, Zain M, Khasrad (2016) The effects of P, S and Mg supplementation of oil palm fronds fermented by *Phanerochaete chrysosporium* on rumen fluid characteristics and microbial protein synthesis. Pakistan Journal of Nutrition 15: 299 – 304. doi: 10.3923/pjn.2016.299.304
2. Febrina D, Jamarun N, Zain M, Khasrad. (2016) Effects of calcium (Ca) and manganese (Mn) supplementation during oil palm frond fermentation by *Phanerochaete chrysosporium* on in vitro digestibility and rumen fluid characteristics. Pakistan Journal of Nutrition 15: 352-358. doi: 10.3923/pjn.2016.352.358.
3. Zahari WM, Hassan OA, Wong HK, Liang JB (2003) Utilization of oil palm frond - based diets for beef and dairy production in Malaysia. Asian-Australasian Journal of Animal Sciences 16 (4): 625 – 634. doi: 10.5713/ajas.2003.625.
4. Rahman MM, Lourenço M, Hassim HA, Baars JJP, Sonnenberg ASM, Cone JW, de Boever J, Fievez V (2011) Improving ruminal degradability of oil palm fronds using white rot fungi. *Animal Feed Science and Technology* 169 (3-4) :157 – 166. doi: 10.1016/j.anifeedsci.2011.06.014.
5. Kawamoto H, Mohamed WZ, Sukur NIM, Ali MSM, Islam Y, Oshio S (2001) Palatability, digestibility and voluntary intake of processed oil palm fronds in cattle. Japan Agricultural Research Quarterly 35; 195 – 200. doi: 10.6090/jarq.35.195.
6. Febrina D, Febriyanti R, Zam SI, Handoko J, Fatah A, Juliantoni J (2018) Antibacterial activity testing and ethanol extract characterization of oil palm fronds (*Elaeis guineensis* Jacq.). Pakistan Journal of Nutrition 17 (9): 427 – 433. doi: 10.3923/pjn.2018.427.433.
7. Sasidharan S, Rajoo N, Rathinam X, Lachimanan YL, Rajoo A (2010) Wound healing potential of *Elaeis guineensis* Jacq. leaves in an infected albino rat model. Molecules 15: 3186 – 3199. doi: 10.3390/molecules15053186.
8. Syahmi, A.R.M.; Vijayarathna, S.; Sasidharan, S.; Latha, L.Y.; Kwan, Y.P.; Lau, Y.L.; Shin, L.N.; Chen, Y (2010). Acute oral toxicity and brine shrimp lethality of *Elaeis guineensis* jacq., (oil palm leaf) methanol extract. Molecules; 15: 8111–8121. doi:10.3390/molecules1511811.
9. Febrina, D (2016) Utilization of oil palm fronds biodelignification products using *Phanerochaete chrysosporium* as substitute forage on goat. Ph.D. Thesis,

- Faculty of Animal Science, University of Andalas, Padang, Indonesia.
10. Franswort, N. R (1996) Biological and Phytochemical Screenings of Plant. *J. Pharm. Sci.*, 1996; 55 (3): 225-265. doi: 10.1002/jps.2600550302.
 11. Handayani, D., N. Sandrawaty, M. Murniati and R. Regina (2015) Screening of endophytic bacteria isolated from marine sponge *Haliclona fascigera* for inhibition against clinical isolates of methicillin resistant *Staphylococcus aureus* (MRSA). *J. Applied Pharmaceut. Sci* 5: 139-142. doi: 10.7324/JAPS.2015.50926.
 12. Ozduven. M. L., F. KOC., C. Polat., L. Coskuntuna (2009) The Effects of Lactic Acid Bacteria and Enzyme Mixture Inoculants on Fermentation and Nutrient Digestibility of Sunflower Silage. *Kafkas Univ Vet Fak Derg* 15 (2): 195-199. doi:10.9775/kvfd.2008.91-A.
 13. Laura. B and G. Paolo (2006) *Algae. Anatomy Biochemistry and Biotechnology*. Boca Raton. CRC Press.
 14. Umiyasih. U dan E. Wina. Processing and Nutritional Value of Corn by-Product as Ruminant Feed. *Wartazoa*. 2008; 18(3):127-136.
 15. Saun RJV, Henrich A.J (Eds) (2008) *Proceedings of the Mid-Atlantic Conference: Pennsylvania, 26 May 2008*. Penn State's Collage.
 16. Amanullah MA, Sekar S, Muthukrishnan P (2010) Prospect and Potential of Poultry Manure. *Asian Journal of Plant Sciences*. 9(4): 1 – 11. doi: 10.3923/ajps.2010.172.182.
 17. Thambirajah JJ, Zulkali MD, Hashim MA (1995) Microbiological and biochemical changes during the composting of oil palm empty-fruit-bunches. Effect of nitrogen supplementation on the substrate. *Bioresource Technology*. 1995; 52(2): 133-144. doi: 10.1016/0960-8524(95)00008-3.
 18. Salamah E, Ayuningrat E, Purwaningsih S (2008) Initial Screening of Fresh Water Mussel Bioactive Components from (*Anodonta woodiana* Lea.) from Antioxidant Compounds. *Buletin Teknologi Hasil Perikanan*.11(2):119-132. doi: 10.17844/jphpi.v11i2.912.
 19. Harborne AJ (1998) *Phytochemical Methods A Guide to Modern Techniques of Plant Analysis*. 3rd Edition. Dordrecht, Springer.
 20. Bose R, Majumdar C, Bhattacharya S (1997) Steroids in *Achatina fulica* (Bowdich): steroid profile in haemolymph and in vitro release of steroids from endogenous precursors by ovotestis and albumen gland. *Comparative Biochemistry and Physiology* 116 (3): 179 – 182. doi: 10.1016/S0742-8413(96)00163-6.
 21. Silva BP, Sousa AC, Silva GM, Mendes TP, Parente JP. A new bioactive steroidal saponin from *Agave attenuata*. *Z Naturforsch* 2002; 57c:423-428.
 22. Schmidt G, Steinhart H (2001) Impact of extraction solvents on steroid contents determined in beef. *Journal of Food Chemistry*. 76:83-88. doi:10.1016/S0308-8146(01)00237-0.
 23. Widya DR, Suryanto D, Desrita (2014) Antimicrobial Activity of Lotus Seeds (*Nymphaea pubescens* L.) on Bacteria *Aeromonas hydrophila*, *Streptococcus*

- agalactiae, and Saprolegnia sp. Jurnal Aquacoastmarine 2(1).7-14.
24. Martono B, Setiyono RT (2014) Phytochemical screening of six tea Genotypes Journal of Industrial and Beverage Crops 1 (2): 63 – 68. doi: [10.21082/jtidp.v1n2.2014.p63-68](https://doi.org/10.21082/jtidp.v1n2.2014.p63-68).
 25. Poelongan M, Praptiwi (2010). Antibacterial activity test of mangos teen (*Gardnia mangostana Linn*) Peel. Media Litbang Kesehatan 20(2):65-69.
 26. Ikalinus R, Widyastuti SK, Setiasih NLE (2015) Phytochemical screening ethanol extract skin stem moringa (*Moringa oleifera*). Indonesia Medicus Veterinus 4(1):71-79.
 27. Wardhani LK, Sulistyani N (2012) Antibacterial activity test of ethyl acetate extract of binahong leaf (*Anredera scandens (L.) Moq.*) against *Shigella flexneri* with the profile of thin layer chromatography. Jurnal Ilmiah Kefarmasian 2(1):1-16.
 28. Nanda A, Saravanan M (2009) Biosynthesis of silver nanoparticles. from *Staphylococcus aureus* and its antimicrobial activity against MRSA and MRSE. Nanomedicine: Nanotechnology, Biology and Medicine 5(4): 452-456. doi:10.1016.j.nano.2009.01.012.
 29. Davis WW, Stout TR (1971), Disc plate method of microbiological antibiotic assay. I. Factors influencing variability and error. Applied Microbiol 22: 659-665.
 30. Mudi SY, Ibrahim H (2008) Activity of *Bryophyllum pinnatum S. Kurz* extracts on respiratory tract pathogenic bacteria. Bayero Journal of Pure Applied Sciences 1: 43-48. doi:10.4314/bajopas.v1i1.57512.

PEMBERITAHUAN DARI EDITOR

“PUBLICATION INVOICE”

15:56

4G+ 45%



[JTLS] INVOICE_Nutritional
Content and Characteristics of
Antimicrobial Compounds

Inbox



Nashi Widodo 1/21/2020


to me



Dear author(s),

We are contacting you in regard to your publication invoice.
You may find
the invoice attached and we appreciate your prompt payment.
Please send your payment proof to this email.

Regards,
Journal of Tropical Life Science
Universitas Brawijaya



Syukria IkhsanZam 1/24/2020

to Journal



Dear JTLS Manager

I hereby submit proof of journal payment



ARTIKEL YANG DIPUBLISH

Research Article

Nutritional Content and Characteristics of Antimicrobial Compounds from Fermented Oil Palm Fronds (*Elaeis guineensis* Jacq.)

Dewi Febrina¹, Rahmi Febriyanti¹, Syukria Ikhsan Zam^{2*}, Zumarni¹, Jepri Juliantoni¹, Abdul Fatah¹

¹ Department of Animal Science, Faculty of Agriculture and Animal Sciences, State Islamic University of Sultan Syarif Kasim Riau, Pekanbaru 28293, Indonesia

² Department of Agrotechnology, Faculty of Agriculture and Animal Sciences, State Islamic University of Sultan Syarif Kasim Riau, Pekanbaru 28293, Indonesia

Article history:

Submission June 2019

Revised September 2019

Accepted November 2019

**Corresponding author:*

E-mail:

syukria.ikhsan.zam@uin-suska.ac.id

ABSTRACT

This research was conducted to determine physical quality, the nutritional quality and characteristics of antimicrobial compounds from fermented oil palm fronds (FOPF). This research was carried out using a Complete Randomized Design with 4 treatments and 4 replications. The following treatments were performed by the addition of different additives namely: 10% poultry manure, 10% cow feces, 5% urea and 5% molasses. All treatments were extracted in stages using n-hexane, ethyl acetate and methanol. The observed parameters were physical quality (pH, aroma, color, and texture), nutritional quality (neutral detergent fiber (NDF), acid detergent fiber (ADF), hemicellulose, cellulose and lignin), yield of extract, class of bioactive compounds and the antimicrobial activity test of extracted compounds. The data were analyzed using a one-way analysis of variance and differences between treatments were tested using Duncan's multiple range test. The addition of different additives has a very significant effect ($P < 0.01$) on the lignin content; significant effect ($P < 0.05$) on pH, aroma, color and texture; and no effect on the content of NDF, ADF, hemicellulose and cellulose. The highest yield of extract was obtained from methanol extract with the addition of poultry manure. The methanol extract on the addition of poultry manure and urea were identified as compounds in the steroid, tannin and phenolic classes. The highest antimicrobial activity test against *Escherichia coli* (12.83 mm) was obtained from FOPF extract (1%) with addition of poultry manure, while highest antimicrobial activity test against *Staphylococcus aureus* (11.67 mm) was obtained from FOPF extract (4%) with addition of molasses. The conclusion of this research was FOPF with addition of poultry manure provides good physical quality, nutritional quality and can inhibit the growth of *E. coli* and *S. aureus* at a concentration of 1%.

Keywords: Nutritional content, Antimicrobial, Oil palm fronds

Introduction

The utilization of oil palm plantation waste such as oil palm fronds as feed is still limited due to the high lignin content [1, 2]. Physical-chemical-biological and combination treatments and the

addition of additives on the fermentation process aim to break down lignocellulose and lignohemicellulose bonds and increase nutrient digestibility [3, 4, 5].

How to cite:

Febrina D, Febriyanti R, Zam SI et al. (2020) Nutritional Content and Characteristics of Antimicrobial Compounds from Fermented Oil Palm Fronds (*Elaeis guineensis* Jacq.). Journal of Tropical Life Science 10 (1): 27 – 33. doi: 10.11594/jtls.10.01.04

Palm oil fronds have antimicrobial activity [6]. Methanol extract from palm oil leaves contains tannin, alkaloids, saponins, terpenoids and flavonoids [7]; and toxicity tests on shrimp and rats showed that palm oil leaf extract is safer for consumption and recommended as a natural product [8]. This study aims to determine physical quality, the nutritional quality, and characteristics of antimicrobial compounds from fermented oil palm fronds.

Materials and Methods

The research was conducted in June-November 2017, at the Laboratory of Nutrition and Chemistry at the Faculty of Agriculture and Animal Sciences of State Islamic University of Sultan Syarif Kasim Riau Indonesia, Biotechnology Research Center of the Indonesian Sciences Institute Indonesia and Laboratory of Biota Sumatra Andalas University.

The study was conducted through two (2) stages: namely fermentation of oil palm frond based on Febrina [9] modified with different additive (10% poultry manure, 10% cow feces, 5% urea and 5% molasses) using a Completely Randomized Design 4 treatment with 4 replications. Fermentation process for 21 days. The data were analyzed using a one-way analysis of variance and differences between treatments were tested using Duncan's Multiple Range Test. The observed parameters were physical quality tests include pH, aroma, color and texture and nutritional quality including the content of NDF, ADF, hemi cellulose, cellulose, and lignin based on Van Soest analysis using the FOSS Fibertec system. The second stage is multilevel maceration and extraction using methanol, ethyl acetate and n-hexane solvents. Phytochemical screening based on Franswort [10] and the antibacterial activity test of oil palm fronds extract based on Handayani *et al.* [11].

Result and Discussion

The Effect of different additives on the physical quality of FOPF

The addition of different additives has a significant effect ($P < 0.05$) on the physical quality of FOPF including pH, aroma, color and texture (Table 1). The lowest of pH (5.18) in treatment A is related to the availability of carbohydrates and the production of lactic acid bacteria. The same results were seen in treatment D with a pH of 5.28. Ad-

dition of poultry manure and molasses on fermentation oil palm frond aims as a source of additive which can trigger the growth of lactic acid bacteria to produce lactic acid. Increased production of lactic acid will reduce pH and inhibit the growth of pathogenic bacteria. Ozduven *et al.* [12] stated that pH is associated with lactic acid produced during the fermentation process which can inhibit the growth of pathogenic microbes. Oil palm fronds which are fermented with molasses produce pH 3.93 [3].

The process of ammonization of oil palm fronds using urea will produce the highest pH of 7.60 which is alkaline. The same results were reported by Zahari *et al.* [3], the process of ammonization of oil palm fronds using urea resulting in a pH of 7.38. In the process of ammonization with urea will break down into ammonium ions so that the ammonization product is alkaline. Laura and Paolo [13] state that urea will break down into ammonium ions, which are alkaline and ammonization processes can break down glycoside bonds [14] so that the oil palm fronds can be degraded by rumen microbes.

In treatment A produces an aroma score of 3 with acid criteria. It has been shown that the fermentation process is going well as indicated by acid production. Aroma is related to the production of lactic acid and pH. It can be seen in Table 1 the addition of poultry manure to oil palm fronds fermentation resulting in the lowest pH (5.18) so that the products are acidic.

Treatment A, B and D produces a color that is not different ($P > 0.05$), which is natural green or yellowish green with score of 3-3.03. Good quality silage will produce colors that are not different from the original material [15]. The occurrence of color changes from green to yellowish green indicated the anaerobic phase oxygen has run out of carbohydrates. Then it will produce CO_2 , H_2O and heat. The heat produced causes a change in color from green to yellowish green.

Treatment A will produce softer textures with score 2. This shows that the addition of the poultry manure to the fermentation process can support the growth of lactic acid bacteria to produce lactic acid. Optimal production of lactic acid will reduce pH so that it inhibits the growth of pathogenic bacteria. This low pH also supports the production of lignase enzymes which function to break down lignocellulose and lignohemicellulose thereby re-

Table 1. The effect of different additives on the physical quality of FOPF

Treatment	Physical quality							
	pH		Aroma		Color		Texture	
	Value	Notes	Value	Notes	Value	Notes	Value	Notes
A	5.18 ^a ±0.10	Acid	3.00 ^b ±0.00	Acid	3.03 ^a ±0.05	Yellowish Green	2.00 ^a ±0.00	Rather Soft
B	5.52 ^b ±0.20	Acid	4.00 ^c ±0.00	More Acidic	3.00 ^a ±0.00	Yellowish Green	3.00 ^b ±0.00	Soft
C	7.60 ^c ±0.14	Base	2.03 ^a ±0.05	Non-acidic	4.03 ^b ±0.05	Brownish Green	3.03 ^b ±0.05	Soft
D	5.28 ^{ab} ±0.21	Acid	5.00 ^d ±0.00	Very Acidic	3.00 ^a ±0.00	Yellowish Green	3.00 ^b ±0.00	Soft

Table 2. The effect of different additive on fiber fraction of FOPF

Treatment	Fiber Fraction (%)				
	NDF	ADF	Hemicellulose	Cellulose	Lignin
A	73.34±3.68	51.81±4.82	21.52±6.07	28.54±3.36	19.94 ^a ±1.47
B	75.39±2.03	55.03±4.23	20.35±4.46	29.57±3.09	23.15 ^b ±0.96
C	73.83±1.01	53.30±3.43	20.51±4.30	31.55±1.99	21.57 ^a ±1.58
D	74.98±3.52	54.04±4.48	20.93±6.63	31.70±4.27	22.41 ^b ±1.22

Note: A = FOPF with the addition of poultry manure

B = FOPF with the addition of cow feces

C = FOPF with the addition of urea

D = FOPF with the addition of molasses

a, b, and c mean with different superscript differ ($p < 0.05$)

ducing lignin content. It is seen that treatment A produced the lowest lignin content of 19.94%. The release of lignin from cellulose and hemicellulose causes fermented products to have a softer texture.

The Effect of different additive on fiber fraction of FOPF

The addition of different additive had no effect ($P > 0.05$) on the content of NDF, ADF, hemicellulose and cellulose but had a very significant effect ($P < 0.01$) on the lignin content (Table 2). Rahman *et al.* [4] reported there was a decrease in the lignin content of oil palm fronds in the fermentation process using white rot fungi.

Lignin is a part of plant cell walls which consists of very complex and difficult to digest bonds. The high content of lignin in the feed will reduce its digestibility. The lignin content of oil palm fronds is 30.18% [2]. One of the purposes of processing feed through ammonization and fermentation in oil palm fronds is to reduce the lignin content. The addition of different additive gave a very significant effect ($P < 0.01$) on the lignin content of FOPF. The lowest lignin content in treatment A (19.94%) was not significantly different ($P >$

0.05) compared to treatment C (21.57%) but was very significant ($P < 0.01$) lower treatment B (23.15%) and treatment D (22.41%).

The lowest lignin content in treatment A (19.94%) allegedly in the oil palm frond fermentation using poultry manure will produce higher lignase and urease enzymes than other treatments. The lignase enzyme is an extracellular enzyme that functions to degrade lignin and urease to break down urea into ammonia. Poultry manure also functions as a source of urease which will break down urea into ammonia and CO₂. The high N content in poultry manure, which is 3.63-5.30% with C/N ratio 5.8-7.6 [16], causes high urease activity so that the addition of poultry manure as a source of additive in palm frond fermentation can reduce lignin content. Thambirajah *et al.* [17] there is a positive correlation between the nitrogen content of poultry manure and the nutritional content of oil palm empty fruit bunches fermented with poultry manure. The addition of 15% poultry manure in ammoniated rice straw can reduce fermentation time from 20 days to 5 days.

Bioactive compounds from extracts of FOPF

Bioactive compounds contained in extracts from FOPF can be known through maceration method with multilevel extraction. This is based on multilevel extraction can separate the bioactive compounds contained in the extract from the FOPF in the same sample based on their level of polarity without the bioactive compounds dissolving in other solvents, therefore in this study the multilevel extraction process starts from a non-polar solvent (n-hexane) followed by a semi-polar solvent (ethyl acetate) and finally with a polar solvent (methanol).

Table 3 shows n acetate extracts FOPF produce the lowest yields ranging from 0.78-1.08% while the methanol extract produces the largest yields ranging from 5.79-7.87%. The same results were reported by Salamah *et al.* [18] the lowest yield of Taiwanese gravestone extract was from n-hexane extract (0.04%) and the highest yield was from methanol extract (0.12%).

FOPF which was extracted with methanol, produced the highest yield of 7.87% of crude compounds. This shows that the most bioactive compounds found in FOPF are polar because they can dissolve in polar solvents namely methanol. Bioactive compounds of FOPF which are non-polar and semi-polar in smaller amounts. Phytochemical analysis of bioactive compounds of extract of fermented oil palm fronds is shown in Table 4.

The extraction process uses methanol solvents in treatments A and C, produce small amounts of tannin and phenolic compounds (+). Sufficient amounts of phenolic compounds (++) were found in treatment B and D. N hexane extract in treatment A produced the lowest yield (0.78%) and methanol extract produced the highest yield (7.87%). This shows that treatment A contains polar compounds (methanol) which are greater than non-polar compounds (n hexane). Analysis of phytochemical compounds shows, treatment A contains bioactive compounds, are steroids and tannins, in small amounts (+) and phenolic compounds in sufficient quantities (++).

a. Steroid

Steroids are initially considered only as components of animal substances (as sex hormones, adrenal hormones, bile acids, etc.), but recently steroids have also been found in plant substances [19]. Steroids contained in methanol extract, ethyl

Table 3. The yield of extracts FOPF

Treatment	Yield of extracts (%)		
	N hexane	Ethyl Acetate	Methanol
A	0.78	1.53	7.87
B	1.08	1.3	7.07
C	1.02	1.17	5.79
D	0.89	1.08	7.4

Note: A = FOPF with the addition of poultry manure
B = FOPF with the addition of cow feces
C = FOPF with the addition of urea
D = FOPF with the addition of molasses
a, b, and c mean with different superscript differ (p < 0.05)

acetate and n hexane in treatment A are thought to be adrenal hormones and sex hormones, because in poultry, metabolic waste that cannot be used again is released as excreta through cloaca. Excreta consists of feces, urine and sperm/egg cells. Bose *et al.* [20] stated that steroid compounds were detected in *Achatina fulica* (freshwater gastropods such as golden snails). Steroids are thought to increase body stamina (aphrodisiac) and anti-inflammation. Silva *et al.* [21] reported extracts from the leaves of *Agave attenuate*, containing steroid compounds that have anti-inflammatory activity.

Phytochemical analysis showed that bioactive steroid compounds were detected in all extract of FOPF (n hexane, ethyl acetate and methanol) which had different polarity. Harbone [19] states, steroids are cholesterol, which are non-polar so they can dissolve in non-polar solvents, (n-hexane), semi-polar (ethyl acetate) and polar (methanol) in small amounts (+). The steroid content in beef, which is extracted using non-polar and polar compounds, shows no difference [22].

Bioactive compounds of steroid, which are found in extract of FOPF can be used as antimicrobials because steroids are bioactive compounds that function as anti-fungal. N hexane extract from lotus seeds contains bioactive steroid compounds and can inhibit the growth of Gram-negative bacteria (*Aeromonas hydrophila*) and Gram-positive bacteria (*Streptococcus agalactiae*) and fungi (*Saprolegnia* sp.) [23].

b. Tannin

Methanol extract from FOPF in treatments A

Table 4. Bioactive compounds of extract of FOPF

Class of Compounds	n hexane				Ethyl Acetate				Methanol			
	A	B	C	D	A	B	C	D	A	B	C	D
Alkaloid	-	-	-	-	-	-	-	-	-	-	-	-
Steroid	+	+	+	+	+	+	+	+	+	+	+	+
Terpenoid	-	-	-	-	-	-	-	-	-	-	-	-
Flavonoid	-	-	-	-	-	-	-	-	-	-	-	-
Saponin	-	-	-	-	-	-	-	-	-	-	-	-
Quinone	-	-	-	-	-	-	-	-	-	-	-	-
Tannin	-	-	-	-	-	-	-	-	+	-	+	-
Coumarin	-	-	-	-	-	-	-	-	-	-	-	-
Phenolic	-	-	-	-	-	-	-	-	+	++	+	++

Table 2. The effect of different additive on fiber fraction of FOPF

Concentration	Inhibited Zone (mm)							
	<i>E. coli</i>				<i>S. aureus</i>			
	A	B	C	D	A	B	C	D
1%	12.83	9	9	7.67	8	7.67	11.33	6.3
2%	11.67	8.67	8.83	7.33	9.16	8.67	10.16	7.67
3%	10.5	8.66	9	7.5	9.67	8	10.16	9.16
4%	8.5	10.33	8	8	7.83	9.33	8	11.67
5%	7.33	9	8.83	7.5	7.83	6.67	9	10

Note: A = FOPF with the addition of poultry manure

B = FOPF with the addition of cow feces

C = FOPF with the addition of urea

D = FOPF with the addition of molasses

- = there are no bioactive compounds

+ = in small quantities

++ = there are in sufficient quantities

and C contains small amounts of tannin compounds (+). The tannin compounds found in oil palm plants are reported by Sasidharan [7] in methanol extract of palm oil leaves; Febrina *et al.* [6] on the ethanol extract of oil palm fronds.

The presence of tannin compounds in methanol extract in treatments A and C shows that methanol extract from FOPF has the potential to be antimicrobial. Tannin has antibacterial properties by precipitation proteins because it is suspected that tannins have the same effect as phenolic compounds [24]. The tannin compounds found in the ethanol extracts of oil palm fronds [7] and mango-steen peel [25] have the potential to be antimicrobial because they can inhibit the growth of *S. aureus* and *E. coli*.

c. Phenolic

Phytochemical analysis in treatment B and D showed that sufficient quantities of phenolic bioactive compounds (++). This indicates that the ex-

tract of oil palm frond which is fermented with cow feces and molasses can be used as an antimicrobial. Phenol compounds can be used as antibacterial because they contain aromatic rings with one or two hydroxyl groups that can change the permeability of the cytoplasmic membrane thus inhibiting bacterial growth [19,26]. Ethyl acetate extract of *binahong* leaves, containing polyphenols and saponins with the minimum inhibitory concentration (MIC) against *Shigella flexneri* was 8% b/v [27].

Antimicrobial activity test of extracts from FOPF (In Vitro)

The methanol extract from oil palm fronds which are fermented with different additive has antimicrobial activity which is characterized by the formation of inhibitory zones. The formation of inhibition zones shows that it has inhibited the growth of Gram-positive bacteria (*S. aureus*) and Gram-negative bacteria (*E. coli*), by compounds

contained in methanol extract from FOPF. The inhibitory zone formed shows that bacteria have antimicrobial activity [28], thus it can be concluded that methanol extract from FOPF which are fermented with different sources of additives has antimicrobial activity that is bactericide (killing bacteria).

The inhibitory zone formed in testing the antimicrobial activity of extracts from FOPF in Gram-positive (*S. aureus*) and Gram-negative (*E. coli*) bacteria at concentrations of 1-5% as shown in Table 5.

In Table 5 shown the difference in diameter of inhibition zones in each treatment. The diameter of the inhibition zone ranges from 6.67 - 12.83 mm. The smallest diameter inhibition zone of 6.67 mm was produced at a concentration of 5% in treatment B. The largest diameter inhibition zone is 12.83 mm at a concentration of 1% in treatment A.

The ability of extracts from FOPF inhibits bacterial growth in the moderate to strong, because the diameter of the inhibition zone ranges from 6.67-12.83 mm. In accordance with the statement of Davis and Stout [29] and Mudi and Ibrahim [30] if the diameter of the inhibition zone is more than 6 mm, the extract has antimicrobial activity with a moderate to strong category.

Table 5 shows the diameter of the inhibitory zone in Gram-negative bacteria (*E. coli*) generally greater than the Gram-positive bacteria (*S. aureus*). The largest diameter of the inhibition zone in Gram-negative bacteria (*E. coli*) namely in treatment A with a concentration of 1% was 12.83 mm. The largest diameter of the inhibition zone in Gram-positive bacteria (*S. aureus*) namely in treatment D with a concentration of 4% is 11.67 mm. There was a difference in the diameter of the inhibitory zone in gram positive and Gram-negative bacteria due to differences in the structure of bacterial cell walls. The data in Table 5 shows that the antibacterial compounds of the methanol extract of FOPF were able to inhibit the growth of Gram-negative bacteria better than Gram-positive bacteria which was characterized by a larger diameter of the inhibition zone. Cell walls of Gram-positive bacteria have a simple structure compared to Gram-negative bacteria that have complex structures. The largest diameter inhibition zone (12.83 mm) for gram negative bacteria (*E. coli*)

showed that the compounds contained in the methanol extract of FOPF had a greater ability to penetrate the cell walls of Gram-negative bacteria. This is because Gram-negative bacteria have a more complex structure and have more layers so that only certain compounds can penetrate it.

The minimum inhibition concentration (MIC) shows the lowest concentration of extract which can inhibit bacterial growth. The smaller the MIC value the greater the antimicrobial activity. This can be seen in treatment A at a concentration of 1% which was able to inhibit the growth of pathogenic microbes with the largest diameter of inhibition zone (12.83 mm).

Conclusion

The conclusion of this research was FOPF with addition of poultry manure provides good nutritional quality and can inhibit the growth of *E. coli* and *S. aureus* at a concentration of 1%.

Acknowledgment

Author would like to thank to the Institute for Research and Community Service of the State Islamic University of Sultan Syarif Kasim Riau which funded this study under contract number 1043/R/2017.

References

1. Febrina D, Jamarun N, Zain M, Khasrad (2016) The effects of P, S and Mg supplementation of oil palm fronds fermented by *Phanerochaete chrysosporium* on rumen fluid characteristics and microbial protein synthesis. *Pakistan Journal of Nutrition* 15: 299 – 304. doi: doi.org/10.3923/pjn.2016.299.304.
2. Febrina D, Jamarun N, Zain M, Khasrad (2016) Effects of calcium (Ca) and manganese (Mn) supplementation during oil palm frond fermentation by *Phanerochaete chrysosporium* on in vitro digestibility and rumen fluid characteristics. *Pakistan Journal of Nutrition* 15: 352 – 358. doi: doi.org/10.3923/pjn.2016.352.358.
3. Zahari WM, Hassan OA, Wong HK, Liang JB (2003) Utilization of oil palm frond - based diets for beef and dairy production in Malaysia. *Asian-Australasian Journal of Animal Sciences* 16 (4): 625 – 634. doi: doi.org/10.5713/ajas.2003.625.
4. Rahman MM, Lourenço M, Hassim HA et al. (2011) Improving ruminal degradability of oil palm fronds using white rot fungi. *Animal Feed Science and Technology* 169 (3-4): 157 – 166. doi: doi.org/10.1016/j.anifeeds.2011.06.014.
5. Kawamoto H, Mohamed WZ, Sukur NIM et al. (2001) Palatability, digestibility and voluntary intake of processed oil palm



- fronds in cattle. *Japan Agricultural Research Quarterly* 35; 195 – 200. doi: doi.org/10.6090/jarq.35.195.
6. Febrina D, Febriyanti R, Zam SI et al. (2018) Antibacterial activity testing and ethanol extract characterization of oil palm fronds (*Elaeis guineensis* Jacq.). *Pakistan Journal of Nutrition* 17 (9): 427 – 433. doi: 10.3923/pjn.2018.427.433.
 7. Sasidharan S, Rajoo N, Rathinam X et al. (2010) Wound healing potential of *Elaeis guineensis* Jacq. leaves in an infected albino rat model. *Molecules* 15: 3186 – 3199. doi: 10.3390/molecules15053186.
 8. Syahmi ARM, Vijayarathna S, Sasidharan S et al. (2010) Acute oral toxicity and brine shrimp lethality of *Elaeis guineensis* Jacq. (oil palm leaf) methanol extract. *Molecules* 15: 8111 – 8121. doi: 10.3390/molecules15118111.
 9. Febrina D (2016) Utilization of oil palm fronds biodelignification products using *Phanerochaete chrysosporium* as substitute forage on goat. PhD thesis. Andalas University, Department Animal Science.
 10. Franswort NR (1996) Biological and phytochemical screenings of plant. *Journal of Pharmaceutical Science* 55 (3): 225 – 265. doi: 10.1002/jps.2600550302.
 11. Handayani D, Sandrawaty N, Mumiaty M, Regina R (2015) Screening of endophytic bacteria isolated from marine sponge *Haliclona fascigera* for inhibition against clinical isolates of methicillin resistant *Staphylococcus aureus* (MRSA). *Journal of Applied Pharmaceutical Science* 5: 139 – 142. doi: 10.7324/JAPS.2015.50926.
 12. Ozduven ML, Koc F, Polat C, Coskuntuna L (2009) The effects of lactic acid bacteria and enzyme mixture inoculants on fermentation and nutrient digestibility of sunflower silage. *Kafkas Universitesi Veteriner Fakultesi Dergisi* 15 (2): 195 – 199. doi: 10.9775/kvfd.2008.91-A.
 13. Laura B, Paolo G (2006) *Algae: Anatomy Biochemistry and Biotechnology*. Boca Raton, CRC Press.
 14. Umiyasih U, Wina E (2008) Processing and nutritional value of corn by-product as ruminant feed. *Wartazoa* 18 (3): 127 – 136.
 15. Saun RJV, Henrich AJ (Eds) (2008) *Proceedings of the Mid-Atlantic Conference: 26 May 2008; Pennsylvania*. Penn States Collage.
 16. Amanullah MA, Sekar S, Muthukrishnan P (2010) Prospect and potential of poultry manure. *Asian Journal of Plant Sciences* 9 (4): 1 – 11. doi: 10.3923/ajps.2010.172.182.
 17. Thambirajah JJ, Zulkali MD, Hashim MA (1995) Microbiological and biochemical changes during the composting of oil palm empty-fruit-bunches. Effect of nitrogen supplementation on the substrate. *Bioresource Technology* 52 (2): 133 – 144. doi: 10.1016/0960-8524(95)00008-3.
 18. Salamah E, Ayuningrat E, Purwaningsih S (2008) Initial screening of freshwater mussel bioactive components from (*Anodonta woodiana* Lea.) from antioxidant compounds. *Buletin Teknologi Hasil Perikanan* 11 (2): 119 – 132. doi: 10.17844/jphpi.v11i2.912.
 19. Harborne AJ. *Phytochemical Methods A Guide to Modern Techniques of Plant Analysis*. 3rd Edition. Dordrecht, Springer.
 20. Bose R, Majumdar C, Bhattacharya S (1997) Steroids in *Achatina fulica* (bowdich): Steroid profile in haemolymph and in vitro release of steroids from endogenous precursors by ovotestis and albumen gland. *Comparative Biochemistry and Physiology* 116 (3): 179 – 182. doi: 10.1016/S0742-8413(96)00163-6.
 21. Silva BP, Sousa AC, Silva GM, Mendes TP, Parente JP. A new bioactive steroidal saponin from *Agave attenuata*. *Zeitschrift für Naturforschung* 57c: 423 – 428.
 22. Schmidt G, Steinhart H (2001) Impact of extraction solvents on steroid contents determined in beef. *Food Chemistry* 76 (1): 83 – 88. doi: 10.1016/S0308-8146(01)00237-0.
 23. Widya DR, Suryanto D, Desrita (2014) Antimicrobial activity of lotus seeds (*Nymphaea pubescens* L.) on bacteria *Aeromonas hydrophila*, *Streptococcus agalactiae*, and *Saprolegnia* sp. *Jurnal Aquacoastmarine* 2 (1): 7 – 14.
 24. Martono B, Setiyono RT (2014) Phytochemical screening of six tea genotypes. *Journal of Industrial and Beverage Crops* 1 (2): 63 – 68. doi: 10.21082/jtidp.v1n2.2014.p63-68.
 25. Poelongan M, Praptiwi (2010) Antibacterial activity test of mangos teen (*Gardnia mangostana* Linn.) peel. *Media Litbang Kesehatan* 20 (2): 65 – 69.
 26. Ikalinus R, Widyastuti SK, Setiasih NLE (2015) Phytochemical screening ethanol extract skin stem moringa (*Moringa oleifera*). *Indonesia Medicus Veterinus* 4 (1): 71 – 79.
 27. Wardhani LK, Sulistyani N (2012) Antibacterial activity test of ethyl acetate extract of binahong leaf (*Anredera scandens* (L.) Moq.) against *Shigella flexneri* with the profile of thin layer chromatography. *Jurnal Ilmiah Kefarmasian* 2 (1): 1 – 16.
 28. Nanda. A, Saravanan M (2009) Biosynthesis of silver nanoparticles from *Staphylococcus aureus* and its antimicrobial activity against MRSA and MRSE. *Nanomedicine: Nanotechnology, Biology and Medicine* 5 (4): 452 – 456. doi: 10.1016/j.nano.2009.01.012.
 29. Davis WW, Stout TR (1971) Disc plate method of microbiological antibiotic assay. I. Factors influencing variability and error. *Applied Microbiology* 22: 659 – 665.
 30. Mudi SY, Ibrahim H (2008) Activity of *Bryophyllum pinnatum* S. Kurz extracts on respiratory tract pathogenic bacteria. *Bayero Journal of Pure Applied Sciences* 1 (1): 43 – 48. doi: 10.4314/bajopas.v1i1.57512.

This page is intentionally left blank.

SCIMAGO JOURNAL RANK (SJR)

“JOURNAL OF TROPICAL LIFE SCIENCE”


Journal of Tropical Life Science







<p>COUNTRY</p> <p>Indonesia</p> <p> Universities and research institutions in Indonesia</p> <p> Media Ranking in Indonesia</p>	<p>SUBJECT AREA AND CATEGORY</p> <p>Agricultural and Biological Sciences</p> <ul style="list-style-type: none"> Ecology, Evolution, Behavior and Systematics Insect Science Plant Science <p>Biochemistry, Genetics and Molecular Biology</p> <ul style="list-style-type: none"> Molecular Biology <p>Environmental Science</p> <ul style="list-style-type: none"> Ecology 	<p>PUBLISHER</p> <p>Brawijaya University</p>	<p>H-INDEX</p> <p>6</p>
<p>PUBLICATION TYPE</p> <p>Journals</p>	<p>ISSN</p> <p>20875517, 25274376</p>	<p>COVERAGE</p> <p>2019-2023</p>	<p>INFORMATION</p> <p>Homepage</p> <p>How to publish in this journal</p> <p>jtls@ub.ac.id</p>

<p>PUBLICATION TYPE</p> <p>Journals</p>	<p>ISSN</p> <p>20875517, 25274376</p>	<p>COVERAGE</p> <p>2019-2023</p>	<p>INFORMATION</p> <p>Homepage</p> <p>How to publish in this journal</p> <p>jtls@ub.ac.id</p>
--	--	---	---

SCOPE

The Journal of Tropical Life [Science](#) (JTLS) provides a rapid publication of full-length Research Papers, Short Communication and Review articles describing new findings or theories in a living system, cells and molecular level in tropical life [sciences](#) and related areas. The journal publishes articles that report novel findings of wide tropical life systems in the areas of biodiversity, agriculture, fisheries, health, **husbandry**, forestry and environmental technology. Research Areas include: • Tropical Forest • Tropical Wildlife • Tropical Environment • Tropical Biomedical Sciences • Tropical Biodiversity • Tropical Bioprospecting • Tropical Ethnobiology • Tropical Diseases • Tropical Bioconservation • Bioinformatics • Biomodeling

 Join the conversation about this journal



Journal of Tropical Life Science
Ecology
Q4
SJR 2023 0.16
best quartile
powered by scimagojr.com

Show this widget in your own website
Just copy the code below and paste within your html code:
`<a href="https://www.scima`

SCImago Graphica
Explore, visually communicate and make sense of data with our **new data visualization tool**.

SCOPUS INDEKS

“JOURNAL OF TROPICAL LIFE SCIENCE”

Source details

Feedback > Compare sources >

Journal of Tropical Life Science

Open Access

Years currently covered by Scopus: from 2019 to 2024

Publisher: Brawijaya University

ISSN: 2087-5517 E-ISSN: 2527-4376

Subject area: Agricultural and Biological Sciences: Insect Science Environmental Science: Ecology Agricultural and Biological Sciences: Plant Science Agricultural and Biological Sciences: Ecology, Evolution, Behavior and Systematics Biochemistry, Genetics and Molecular Biology: Molecular Biology

Source type: Journal

View all documents > Set document alert Save to source list

CiteScore 2023	1.0
SJR 2023	0.155
SNIP 2023	0.356

CiteScore CiteScore rank & trend Scopus content coverage

CiteScore 2023

$$1.0 = \frac{174 \text{ Citations 2020 - 2023}}{178 \text{ Documents 2020 - 2023}}$$

Calculated on 05 May, 2024

CiteScoreTracker 2024

$$1.0 = \frac{180 \text{ Citations to date}}{188 \text{ Documents to date}}$$

Last updated on 05 November, 2024 • Updated monthly

CiteScore CiteScore rank & trend Scopus content coverage

CiteScore 2023

$$1.0 = \frac{174 \text{ Citations 2020 - 2023}}{178 \text{ Documents 2020 - 2023}}$$

Calculated on 05 May, 2024

CiteScoreTracker 2024

$$1.0 = \frac{180 \text{ Citations to date}}{188 \text{ Documents to date}}$$

Last updated on 05 November, 2024 • Updated monthly

CiteScore rank 2023

Category	Rank	Percentile
Agricultural and Biological Sciences		
Insect Science	#143/181	21st
Environmental Science		
Ecology	#368/461	20th
Agricultural and Biological Sciences		
	#433/516	16th

View CiteScore methodology > CiteScore FAQ > Add CiteScore to your site >