



ICSAE-7

**7th International Conference
on Sustainable Agriculture and Environment**

CONFERENCE GUIDE AND BOOK OF ABSTRACTS

**7th International
Conference
on Sustainable Agriculture
and Environment**

August 27, 2020
Surakarta - Indonesia

THE 7th INTERNATIONAL CONFERENCE ON SUSTAINABLE AGRICULTURE AND ENVIRONMENT (ICSAE-7)

Conference Guide and Book of Abstracts

SURAKARTA - INDONESIA

AUGUST 27, 2020

Table of Contents

| | |
|--------------------------------|----|
| Preface | 3 |
| Organizing Committee | 4 |
| Conference Guide | 6 |
| Parallel Session Guide | 7 |
| General Assembly Rundown | 8 |
| Pararell room | |
| Poster Room 1 | 9 |
| Poster Room 2 | 11 |
| Poster Room 3 | 12 |
| Video Room 1 | 13 |
| Video Room 2 | 15 |
| Video Room 3 | 16 |
| Video Room 4 | 17 |
| Video Room 5 | 18 |
| Video Room 6 | 19 |
| List of Abstracts | 20 |

Preface

Contemporary events shows us that we humans are very vulnerable to economic and environmental fluctuations. As humanity progress, produce and consume like it will exist until infinity. Sustainability has always been the case, is accepted as life guidance but until then we; researchers, companies and students should try to work on this concept so that next generation can benefit from our efforts.

Sustainable way of dealing with environment and agriculture is not easy. The benefits from sustainable agriculture vary and sustainable agriculture doesn't have harmful effects on environment or sources. In agricultural activities have been performed for chaining the environment or utilizations of environment as sources with maximum benefits. If we do not success that, it is impossible for us to talk about modern agriculture. Under poor management of agriculture, biodiversity and soil properties can be damage, contaminate and some problems limiting efficient agricultural activities may be occur.

The main objective of farming is to produce food for human being, but unfortunately some parts of the farmlands have used also as settlements areas. By this way, some natural resources have been deteriorated; consequently size of farmland has reduced day by day. In that regard, it is possible to say that we are in going in a circle. More than one-third of food supplies is as form of wastes due to the over buying than required amounts of food by costumers.

In the lights of the information mentioned above, there is no doubt that continues sustainable agricultural activities are practical solution to prevent the human being from the starvation. The International Conference on Sustainable Agriculture and Environment (ICSAE) series were aimed to provide a platform for researchers and academics as well as practicing professionals from all over the world, to present their research and professional development activities in agriculture, environment, food and other relevant subjects. This conference series was an effort to identify the ideas, practices and policies that constitute our concept of sustainable agriculture. The concept of sustainable agriculture itself is still evolving and thus, we published not as a definitive or final statement, but as an invitation to continue our dialogue. Moreover, as we experinced lately during COVID-19 pandemic situation, agriculture is a sector which stay struggle and can be souce of many materials for fight the diseases.

Now, I would like to thank whole people being at Scientific Board, Managerial Board, authors or participants by supporting their valuable works and particularly Prof. Ahmad Yunus from Universitas Sebelas Maret Indonesia and his team, for all their contributions on continuity of these conference series since it was held, for the 1st, at 2013 in Surakarta Indonesia, then move around the globe and at 7th ICSAE its back again in Surakarta. I do appreciate all the efforts given to prepare this event. This is the first digital conference of ICSAE series which could be reach wider and bigger participant around the world.

ICSAE-7 Chair

Organizing Committee

Honorary Planning Committee (in alphabetical order)

1. Prof. Dr. Ahmad Yunus, Vice Rector 1, Sebelas Maret University, Indonesia
2. Dr. Mustafa Şahin, President, Selcuk University, Turkey
3. Dr. Muhammad Sarjan, Mataram University, Indonesia
4. Prof. Dr. Samanhudi, Dean Faculty of Agriculture Sebelas Maret University, Indonesia

Conference Chairs (in alphabetical order)

1. Dr. Mithat Direk, Agricultural Economy, Selcuk University, Konya, Turkey
2. Dr. Sigist Prastowo, Animal Science, Sebelas Maret University, Indonesia

Technical Program Committee (in alphabetical order)

1. Dr. Ahmad Mohammed Ahmed, Agribusiness & Applied Economic, Tanta University, Egypt
2. Dr. Azam Kakar, Biotechnology, Quetta, Pakistan

Conference Secretary (in alphabetical order)

1. Dr. Adi Ratriyanto, Animal Science, Sebelas Maret University, Indonesia
2. Dr. Amalia Tetrani Saky, MSc. Department of Agrotechnology, Sebelas Maret University, Indonesia

Treasurer

1. Prof. Dr. Sri Hartati, Agrotechnology, Sebelas Maret University, Indonesia

Organizing Committee Members (in alphabetical order)

1. Ms. Dessya Putri
2. Ms. Rina Puji Lestari
3. Ms. Rissa Rahmadwiyati
4. Mr. Rohmad Setiaji
5. Ms. Wafa Nur Hanifah
6. Ms. Winna Candra M

Scientific Committee Members (in alphabetical order)

1. Prof. Dr. Ahmed Mohamed Ahmed, Tanta University, Egypt
2. Dr. Aluh Nikmatullah, Agroecotechnology, Faculty of Agriculture, Mataram University, Indonesia
3. Dr. Anissa Gara, Institut national agronomique de Tunisie and Institut national de recherches agronomiques de Tunisie, Tunisia
4. Dr. Asghar Ali, Institute of Agricultural and Resource Economics, Faculty of Social Sciences, University of Agriculture, Pakistan
5. Prof. Dr. Bambang Hari Kusumo, Postgraduated Program, Mataram University, Indonesia
6. Prof. Dr. Bambang Pujiasmanto, College of Agriculture, Sebelas Maret University, Indonesia
7. Dr. Bilal Acar, Irrigation and Agricultural Building Department, Selçuk University, Turkey

8. Dr. Bilal Cemek, Agricultural & Biosystems Engineering, Ondokuz Mayıs University, Turkey
9. Prof. Dr. Edi Purwanto, Agrotechnology, Sebelas Maret University, Indonesia
10. Dr. Eka Handayanta, Animal Science, Sebelas Maret University, Indonesia
11. Associate Prof. Dr. Hela Chikh Rouhou, Centre Régional des Recherches en Horticulture et Agriculture Biologique (CRRHAB/IRESA), Tunisia
12. Luthfi Ahmaddani, IPB University, Indonesia
13. Prof.Dr. Midhat Jazic, Univerzitet U Tuzli Tehnološki Fakultet Tuzla, Federation of Bosnia And Herzegovina
14. Prof. Dr. Muhammad Ashfaq, Institute of Agricultural and Resource Economics, Faculty of Social Sciences, University of Agriculture, Pakistan
15. Prof. Dr. M. Hasil Tamzil, Animal Science Faculty, Mataram University, Indonesia
16. Dr. Muhammad Khalid Bashir, Institute of Agricultural and Resource Economics, Faculty of Social Sciences, University of Agriculture, Pakistan
17. Prof.Dr. Ovais Omer, University of Veterinary and Animal Sciences, Pakistan
18. Dr. Prabang Setyono, Environmental Science, Sebelas Maret University, Indonesia
19. Assistant Professor Dr. Prithwiraj Jha, Raiganj Surendranath College, India
20. Dr. Şenay Aydın, Department of Soil Sciences, Celal Bayar University, Turkey
21. Dr. Siti Latifah, Forestry Program Study, Mataram University, Indonesia
22. Prof. Dr. Venti Suryanti, Faculty of Math and Science, Sebelas Maret University, Indonesia
23. Dr. Wayan Swana, Faculty of Mathematic and Natural Science, Mataram University, Indonesia
24. Dr. Zubia Masood, SBK Women's University, Pakistan

Video Room 4

| THEME : NANOTECHNOLOGY AND PRECISION FARMING | | |
|--|--|----------|
| Authors | Title | Paper ID |
| Yulensri, Misfit Putrina dan Kresna Murti | Applications of bio-liquid organic fertilizer (bio-lof) to increase the growth and production of corn (<i>Zea mayz</i> L.) | 25 |
| Atika Ramadhanty Puteri, Nanik Setyowati, Fahrurrozi dan Zainal Mukhtar | Growth and Yield of Sweet Corn (<i>Zea Mays</i> L. Saccharata) as Affected by Incubation Time of Preparation for Tithonia (<i>Tithonia diversifolia</i>) Enriched Liquid Organic Fertilizer | 27 |
| Y Effendi, A El Hadi, A Pambudi | Consortium bacteria design of pathogenic fungi development controller fusarium <i>Oxysporum f. sp. Cubense</i> (foc) | 45 |
| S Denaya, R Yulianti, A Pambudi, Y Effendi | Novel microbial consortium formulation as Plant Growth Promoting Bacteria (PGPB) agent | 46 |
| Restu Puji Mumpuni, Ariika Jamilatur Rohmah | Application of biological control <i>Paenibacillus polimixa</i> toward bacterial leaf blight disease on rice plant | 59 |
| Abdul Kadir Salam, M. Afif Hidayatullah, and Supriatin, and Sri Yusnaini | The phytoextraction of cu and zn by elephant grass (<i>Pennisetum purpureum</i>) from tropical soil 21 years after amendment with industrial waste containing heavy metals | 63 |
| Nilahayati | Initial growth of kijas putih soybean (<i>Glycine max</i> (L) merril) in various dosage of gamma rays irradiation | 77 |
| Juwartina Ida Royani, Widia Sri Fatma, Susiyanti, Andi Apriany Fatmawaty | The effect of benzilaminopurine and kinetin for shoot multiplication of indigofera (<i>Indigofera zollingeriana</i> miq.) By in vitro culture | 78 |
| Sabila Awanis, Edi Purwanto, Parjanto | Morphophysiological characteristics and yield of black rice (<i>Oryza sativa</i> L. Indica) induced by gamma rays 100 Gy at M3 generation | 114 |
| Salma S, Elsanti, LN Neneng, A Dariah, Husnain | Biodecomposer of seaweed composting | 116 |
| Yonny Koentjoro, Sukendah, Edi Purwanto, Djoko Purnomo | The role of silicon on content of proline, protein and abscisic acid on soybean under drought stress | 132 |
| Wayan Wangiyana, I Gusti Putu Muliarta Aryana, Ni Wayan Dwiani Dulur | Mycorrhiza biofertilizer and intercropping with soybean increase anthocyanin contents and yield of upland red rice under aerobic irrigation systems | 133 |
| Ramdan Hidayat ^{1*} , Maharani Putri Puji Artiningrum ² , Pangesti Nugrahani | Study of planting media and nutrition concentration on growth rate and yield of Lettuce (<i>Lactuca sativa</i> L.) in NFT hydroponic systems | 141 |

PAPER • OPEN ACCESS

Germination and seedling growth of Kipas Putih soybean (*Glycine max* [L.] Merrill) in various dosage of gamma rays irradiation

To cite this article: Nilahayati *et al* 2021 *IOP Conf. Ser.: Earth Environ. Sci.* **637** 012052

View the [article online](#) for updates and enhancements.

Germination and seedling growth of Kipas Putih soybean (*Glycine max* [L.] Merrill) in various dosage of gamma rays irradiation

Nilahayati¹, D S Hanafiah² and Rosmaina³

¹Department of Agroecotechnology, Faculty of Agriculture, Universitas Malikussaleh, Aceh 24354, Indonesia

²Department of Agroecotechnology, Faculty of Agriculture, Universitas Sumatera Utara, Medan 20155, Indonesia

³Department of Agrotechnology, Faculty of Agriculture and Animal Science, Universitas Islam Negeri Sultan Syarif Kasim Riau, Panam Campus-Pekan Baru 28293, Riau, Indonesia

Corresponding author: nilahayati@unimal.ac.id

Abstract. Gamma irradiation is the widely used mutagen in assembling plant genetic diversity. The important thing to do is looking for the most appropriate dose that gives rise of diversity without turning off irradiated plants. This study aims to determine the initial growth response of Kipas Putih soybean seeds due to gamma ray irradiation treatment. Kipas Putih soybean seeds measure their water content up to 11%. Furthermore, gamma ray irradiation was carried out at doses 0, 200, 400, 600, 800 and 1000 Gray. The results showed that there was decrease percentage of germination in Kipas Putih soybean seeds at 200 Gray reaching 50% at 14th day after planting. Irradiation doses of 400 Gray and 600 Gray reduce percentage of germination until 20% and 30%, respectively. While at doses of 800 Gray and 1000 Gray, all seeds died at 21st day after planting. Giving gamma ray irradiation also showed a decrease in seedling height along with increasing irradiation dose.

1. Introduction

Soybean (*Glycine max* [L.] Merrill) is a very important and miraculous plant in the 21st century. Soybean is in great demand because it has benefits in fulfilling human foods and animal feed. According to Malek, et al., [1] soybean contains vegetable protein which is rich in essential amino acids. Soybean oil also fulfills 30% of the world's vegetable oil which is friendly for people with heart disease. Furthermore, soybeans contain lactose free fatty acids, antioxidants and folic acids, vitamin B complex and isoflavones.

The increasing need of soybeans must be followed by increased production. Various efforts can be taken to achieve this goal. One of them is by assembling new high yielding varieties. A superior variety assembly program can be carried out using plant breeding techniques. The genetic diversity of population is the basis for starting a plant breeding program by plant breeder. Broad genetic diversity can be the basis for successful genetic improvement through selection in plant breeding. There will be no improvement in heritable plant character without genetic diversity. This happens



because if a character has narrow genetic diversity or variability, then each individual in the population is almost the same, making it difficult to improve character through selection.

Mutation breeding is one of technology in increasing diversity to improve the desired plant characteristics. Manjaya et al., [2] stated that mutation breeding is a way to increase the genetic variability of plant characters, both in quantitative and in qualitative manners. The important benefit of mutation breeding is that mutants with certain characters can be created without changing the essential character of the existing variety. Therefore, the mutants can be easily cultivated as a normal type.

Gamma rays are the most energetic form of electromagnetic radiation. Their energy level is from ten to several hundred kilo electron volts. Gamma rays are considered as the most penetrating compared to other radiations [3]. Gamma rays can cause changes at both the gene and chromosome level. Changes in genes and chromosomes will result in changes in plant characters. It is expected to increase genetic diversity in the Kipas Putih variety.

Hanafiah et al., [4] reported gamma ray irradiation treatment to Argomulyo soybeans at 200 Gray, 400 Gray, 600 Gray, 800 Gray and 1000 Gray. The results showed that the lethal dose (LD 50) was at a dose of 457.136 Gray. The IAEA (International Atomic Energy Agency) recommended gamma ray irradiation dosage for soybean plants is 200 Gray, which is useful for improving the quantitative character of plants. The irradiation dose is different for each cultivar and soybean varieties [5] Therefore, it is necessary to conduct research to determine irradiation dose for the Kipas Putih variety which can induce the highest genetic diversity without causing plant mortality. This is an early stage research which aims to find irradiation doses that will be used to obtain mutant lines of soybean plants which can later be released as high yielding and early maturing varieties.

2. Materials and methods

Soybean seed irradiation is carried out at the Center for Isotope and Radiation Technology Research and Development, National Nuclear Energy Agency (BATAN), Jakarta. The irradiation doses used were 0 Gray, 200 Gray, 400 Gray, 600 Gray, 800 Gray and 1000 Gray, and the dose rate was 0.96481 kgray hour⁻¹ (96.481 krad hour⁻¹). A dose of 0 Gy was carried out as a control treatment. This research used Kipas Putih soybean seeds (water content 11%).

After being irradiated with a dose according to each treatment, soybean seeds were planted in polybags for seed germination. The medium for the germination of polybags was soil, sand and manure with 1: 1: 1 ratio. Each treatment contained 30 seeds so that the total number of seeds was 180 seeds. Watering was done in the morning or afternoon in accordance with the conditions to keep the soil moist.

The observation was made as percentage of germination in 2 weeks after planting and seedling height. The calculation is done by dividing the total number of seeds that grow after being irradiated by all the seeds irradiated with gamma rays then multiplying by 100%. The seedling height is measured from the base of the stem to the tip of the seed. Seedling height measurements were carried out at 7th, 14th and 21st days after germination.

3. Results and discussion

The results showed the decrease in the percentage of germination by giving gamma ray irradiation to soybean seeds compared to the control. The germination percentage ranges from 60% (200 Gray) to 96.6% (0 Gray). The treatment of gamma irradiation at 200-1000 Gray showed 60 -76.6% germination percentage at 14th day after planting (Table 1). The damage due to mutagenic substances in the early stages of cell division at high irradiation doses affects seed germination. The embryos in the seeds were subjected to mutagen treatment.

Present result are in accordance with the finding of Ramesh et al., [6] in rice, [7] in cowpea, [8] in mungbean, [9] in bhendi, [10] in rosella and [11] in soybean. [12] also reported a decrease in germination of soybean seeds given gamma ray mutagen on Denna 1 soybean. They irradiated soybean seeds with ⁶⁰Co gamma rays at doses of 0, 100, 200, 300, and 400 Gray. The results showed that the highest percentage of seed germination at 3 days after planting was found at 100 Gray (74%),

while the lowest was at 200 Gray and 400 Gray, namely 67.7%. Similar results were also found by [13] in soybean. He found the different doses of gamma rays treatment showed gradual reduction of seed germination (10thday), seedling survival (30thday) and plant height (30thday) than control in soybean seed var. Co-1. The effects of gamma rays were different doses on survival percentage, mutation frequency and mutagenic effectiveness. The survival percentage and mean value of M₁ generation were decreased with increase the dose of treatments.

Table 1. The percentage of germination due to various dosage of gamma ray irradiation in Kipas Putih soybean seed.

| Doses (Gray) | Germination (%) | | | Average (%) |
|--------------|-----------------|-----|----|-------------|
| | 1 | 2 | 3 | |
| 0 | 100 | 100 | 90 | 96.66 |
| 200 | 50 | 60 | 70 | 60 |
| 400 | 80 | 50 | 70 | 66.66 |
| 600 | 70 | 70 | 60 | 66.66 |
| 800 | 70 | 70 | 60 | 66.66 |
| 1000 | 80 | 90 | 60 | 76.66 |

The decline number of survival plants can be seen clearly in this study. The plants still survive even though only 30 -50 % at 200 - 400 Gray doses. In the mutagen treatment at a dose of 600-1000 Gray, most plants were only able to germinate until they released cotyledon leaves, no growth occurred at the next stage. One month after planting, all plants in this treatment died (Figure 1).



Figure 1. Effects of gamma rays on the seedling growth of Kipas Putih Soybean at 14th day after germination

The maximum reduction was recorded in higher doses of gamma rays (1000 Gray). This was in collaboration with the earlier reports [4]. they found the increased of irradiation dose up to 800 Gray, there was a decrease in germination. Whereas, the germination continued at 1000 Gray, but the sprouts growth stagnation and have abnormalities. The stems and cotyledon were thickened, and the plant height remained (not increased) until the end of observation (two weeks after planting). This was in confirmation with the finding of [14] in bhendi. The results showed that there was a decrease in the number of survival plants treated with gamma irradiation compared to control plants. This was also reported by Satpute et al., [15]The result indicated that lower doses of the mutagen are more effective for induction and recovery of mutations for improvement of soybean.




Figure 2. Seedling height of Kipas Putih soybean due to gamma ray irradiation at 7th, 14th and 21st day after germination

The present study revealed that the seedling height decreased progressively as the doses of mutagens treatment increased at 7th, 14th and 21st days after planting (Figure 2). Seedling height was maximum in control than the mutagenic treatment. The reduction in seedling height on 21st day varied from 47.4 cm (0 Gray) to 2.2 and 2.6 cm (800 Gray, 1000 Gray). The highest seedling height reduction was observed at 600, 800 and 1000 Gray treatment. There are many early reports on dose depended reduction on seedling height in soybean given by [16,17]. A reduction in seedling height can be attributed to the inhibition of growth due to low rate of cell division, decreased amylase activity and increased peroxide activity.




Figure 3. Effects of gamma rays on the seedling growth of Soybean (*Glycine max* L.) cv. Kipas Putih A. at 7th day after planting; B. at the 14th day after planting

The control plants (0 Gray) showed the growth of cotyledons and plumula leaves which had opened perfectly (Figure 3). The 200 Gray irradiation dose treatment showed the growth of cotyledon leaves which had opened completely but the plumule leaves had not appeared. Plants with a dose of 400-1000 Gray showed that many cotyledon leaves had not opened completely and the plumule leaves had not yet appeared. The higher the irradiation dose, the lower the seedling height. At a dose of 600-1000 Gray, it was also seen that the plants could only germinate until they sprouted the cotyledon leaves. Furthermore, the longer it does not show any further growth.

Jagajantham et al. [14] noted that the biological effect of gamma ray was determined by the interaction among atoms or molecules in the cell. These radicals could damage or modify the important components of plant cells, and changed morphology, anatomy, biochemistry and physiology

of plants depending on the radiation dose. The irradiation of seeds with high doses of gamma ray disturbs the synthesis of protein, hormone balance, leaf gas exchange, water exchange and enzyme activity [18]. Girija and Dhanavel [19] stated the ability of mutagens to enter the cells of living organisms for interacting with the DNA produces toxic that associate with their mutagenic properties. Thus, mutagens can cause physiological damages mainly showed by growth retardation and death in M₁ generation.

4. Conclusion

Gamma ray irradiation in various dosage affects percentage of germination and seedling height of Kipas Putih soybean seed. The dosage under 400 Gray can be used to induce genetic diversity in Kipas Putih soybean.

References

- [1] Malek M A, Rafii M Y, Afroz M S S, Nath and Mondal M M A 2014 *Sci. World J.* 1–2
- [2] Manjaya J G and Nandanwar R S 2007 *Plant Mutat. Reports* **1** 36–40
- [3] Ramya B, Nallathambi G and Ram G 2014 *African J. Biotechnol.* **13** 951–6
- [4] Hanafiah D S, Trikoesoemaningtyas, Yahya S and Wirnas D 2010 *Biosfera* **27** 103–11
- [5] Srisombun S, Benjamas K, Chitima Y and Jeeraporn K 2009 *Soybean variety improvement for high grain protein content using induced mutation (Vietnam)*
- [6] Ramesh D V and Seetharami T V V R 2002 *Cytol. Genet* **3** 115–20
- [7] Gnanamurthy, Mariyammal S, Dhanavel D and Bharathi T 2012 *Inter. J. Res. Plant Sci.* **2** 39–42
- [8] Tah R S 2006 *Asian J. Plant Sci.* **5** 61–70
- [9] Elangovan R and Pavadai P 2015 *Hortic. Biotechnol. Res.* **1** 35–8
- [10] Hanafiah D S, Siregar L A M and Putri M D 2017 *Int. J. Agric. Res.* **12** 28–35
- [11] Pavadai P, Girija M and Dhanavel D 2010 *J. Eco. Bio. Tech* **2** 47–50
- [12] Harsanti L and Yulidar 2015 *Prosiding dan presentasi ilmiah. Penelitian Dasar Ilmu Pengetahuan dan Teknologi Nuklir* pp 59–63
- [13] Pavadai P 2015 *Int. J. Mod. Cell. Mol. Biol.* **4** 1–10
- [14] Jagajanantham N, Dhanavel D, Pavadai P and Chidambaram A A 2012 *Int. J. Res. Plant Sci.* 56–8
- [15] Satpute R A and Fultambkar R V 2012 *Curr. Bot.* **3** 18–20
- [16] Pepol P and Pepo P 1989 *Soybean Abstr.* **12** 4–7
- [17] Kusmiyati F, Sutarno M G A S and Herwibawa B 2017 *IOP Conf. Series: Earth and Environmental Science* p 012059
- [18] Toker C, Uzun B, Canci H and Ceylan F O 2005 *Rad Phy Chem* **73** 365–7
- [19] Girija M and Dhanavel D 2013 *Int. J. Res. Biol. Sci.* **3** 84–7



CERTIFICATE

THIS IS TO CERTIFY THAT

Dr. Rosmaina, S.P, M.Si

HAS PARTICIPATED AS

PARTICIPANT

*at the 7th International Conference
on Sustainable Agriculture and Environment (ICSAE-7)*

*August 27, 2020
Surakarta - Indonesia*

Vice Rector of Academic Affairs



Prof. Dr. Ir. Ahmad Yunus, MS
Universitas Sebelas Maret - Indonesia

Chairman of ICSAE



Dr. Mithat Direk
Selçuk University - Turkey