



Restu Misrianti <misriantirestu@apps.ipb.ac.id>

[BuletinPeternak] Submission Acknowledgement

1 message

Buletin Peternakan <buletin.fapet@ugm.ac.id>
To: Restu Misrianti <misriantirestu@apps.ipb.ac.id>

Tue, Jun 22, 2021 at 9:09 AM

Restu Misrianti:

Thank you for submitting the manuscript, "Determination of Morphological Characteristic in Kuantan Cattle Using Multivariate Analysis" to Buletin Peternakan. With the online journal management system that we are using, you will be able to track its progress through the editorial process by logging in to the journal web site:

Manuscript URL:

<https://journal.ugm.ac.id/buletinpeternakan/author/submission/66868>

Username: restumisrianti

If you have any questions, please contact me. Thank you for considering this journal as a venue for your work.

Buletin Peternakan

Buletin Peternakan

=====Email Signature=====

This email was sent automaticly by Buletin Peternakan system, If you have any questions, please contact me :

Buletin Peternakan, Bulletin of Animal Science

Fakultas Peternakan UGM, Jl. Fauna No.3, Bulaksumur, Kampus UGM, Yogyakarta, Indonesia, 55281

Phone Hunting : +62 857 0125 1919 (Sri Agtin Tejawati)

Fax: +62 274 521578

Email : buletin.fapet@ugm.ac.idWebsite : <http://buletinpeternakan.fapet.ugm.ac.id> or<https://journal.ugm.ac.id/buletinpeternakan/index>Twitter : <https://twitter.com/buletinpeternak>

ISSN 0126-4400 (Print)

ISSN 2407-876X (Online)

Prefix DOI : 10.21059

Buletin Peternakan has been accredited as a scientific journal by the Ministry of Research-Technology and Higher Education Republic of Indonesia: 36a/E/KPT/2016 PERIOD : 2016-2021.

Buletin Peternakan published by :

Faculty of Animal Science, Universitas Gadjah Mada

and

ISSTAP (Indonesian Society for Sustainable Tropical Animal Production)



Restu Misrianti <misriantirestu@apps.ipb.ac.id>

[BuletinPeternakan] Editor Decision

2 messages

Buletin Fapet <buletin.fapet@ugm.ac.id>
To: misriantirestu@apps.ipb.ac.id, jakaria@apps.ipb.ac.id

Sun, Aug 8, 2021 at 8:43 AM

Jakaria:

Naskah saudara yang berjudul "Determination of Morphological Characteristic in Kuantan Cattle Using Multivariate Analysis". Dapat dipertimbangkan untuk diterima dengan perbaikan sesuai dengan saran reviewer dan editor, naskah hasil perbaikan akan Kami tinjau kembali. Berikut Kami lampirkan naskah koreksi dari reviewer dan editor beserta komentarnya. Perubahan dalam teks harap diberi warna yang berbeda. Kami hanya menerima 1 file hasil revisi dari penulis. Kami harapkan menerima naskah revisi dari penulis maksimal 7 hari setelah menerima email ini yaitu 15 Agustus 2021.

Our decision is: Revisions Required

Comment Reviewer A:
Dapat dilihat pada lampiran

Comment Reviewer B:
The manuscript great information and writing with correct grammatically

Comment Editor:
Naskah dapat diterima, perbaiki sesuai saran dari reviewer

Official Buletin Peternakan
Faculty of Animal Science, Universitas Gadjah Mada, Indonesia
buletin.fapet@ugm.ac.id

2 attachments

 **Reviewer A.docx**
4304K

 **Reviewer B.docx**
6728K

Restu Misrianti <misriantirestu@apps.ipb.ac.id>
To: Buletin Fapet <buletin.fapet@ugm.ac.id>

Mon, Aug 9, 2021 at 8:28 PM

Kepada yth Dewan Editor
Buletin Peternakan Fapet UGM

Terima kasih saya ucapkan atas saran dan masukan untuk naskah saya yang berjudul "Determination of Morphological Characteristic in Kuantan Cattle Using Multivariate Analysis". Berikut ini saya lampirkan naskah yang sudah saya perbaiki sesuai masukan dan saran dari reviewer dan editor (Perubahan dalam teks, saya tandai dengan warna kuning). Demikian saya sampaikan, atas perhatian Bapak dan Ibu, saya ucapkan banyak Terima Kasih.

Wassalam
Restu Misrianti
[Quoted text hidden]

 **Determination of Morphological Characteristic in Kuantan Cattle Using Multivariate Analysis_08082021.docx**
6871K

1 **Determination of Morphological Characteristic in Kuantan Cattle Using**
2 **Multivariate Analysis**
3
4

ABSTRACT

The objective of this research was to characterized morphology and estimated genetic distance between the population of Kuantan cattle. A Total of 213 cattle (44 male and 169 female with age ranging from 2-3 years) were used in this study and collected from extensive ranging systems in three sub-population (Cerenti, Inuman, and Kuantan Hilir regions) Kuantan Singingi Regency, Riau Province. Five variables were measured that is Body Length (BL)(cm), Withers Height (WH)(cm), Hip Height (HH)(cm), Chest Girth (CG)(cm), and Chest Depth (CD)(cm). Data obtained were analyzed, descriptive analysis, Principal Components Analysis (PCA) and Hierarchical Clustering Analysis (HCA) using XLSTAT program. All variables of body measurement in the Kuantan Hilir region were higher than Cerenti dan Inuman, Kuantan Singingi Regency, Riau Province. The first factor in PCA described body measurement contributed 32.77%, and the second factor described body shape contribute 25.83% of total variability. The dendrogram showed there is three clusters of Kuantan Cattle.

Keywords: Kuantan cattle, PCA, Genetic Distance, Morphological Characteristics

Introduction

Indonesia has a large variety of indigenous animal genetic resources (AnGR). One of them is Kuantan cattle. Kuantan cattle are one of the local breeds in Riau Province (Indonesia) that play an important role in the maintenance of rural population living, social, religious, and traditional celebration. Kuantan cattle in Riau Province very adaptive in an extensive management system, and lack quality feeder. Based on data in the Kuantan Singingi government, the population of Kuantan cattle decreasing from 2016/2017.

Commented [A1]: Tampilkan data populasi dan penurunannya sejak 2016/2017 hingga sekarang.

1 90The decrease in grazing areas and lack of bulls make the population decrease.
2 The absence of a structured and targeted breeding program in Kuantan cattle, and the
3 crossing of Kuantan cattle with other breeds without evaluation, is feared genetic drift.
4 FAO (201P5) states that cattle are categorized into endangered groups if the total
5 number of males is less than or equal to 20 and greater than 5.

6 Conservation and production program require strategy in management and detail
7 information about animal genetic resources (FAO, 2013). Morphological characterization
8 of the breed, very important to develop the breeding scheme. Improvement in the
9 breeding system is very important to increase the population and to improve the quality
10 of Kuantan cattle.

11 The multivariate technique is one of the methods to characterize local breed and
12 has been found very suitable to identified genetic variation within and between
13 populations. PCA can identify the major quantitative variable, and discriminant analyses
14 more suitable in assessing to evaluate their differentiation. Recently, this technique has
15 been suitable to characterize Indonesia local breed cattle (Bali cattle (Hikmawaty *et al.*
16 2017), Pasundan Cattle (Sulasmı *et al.*, 2017)), Ongol and Sumba Ongol (Putra *et al.*,
17 2020) Aceh Cattle (Mahmudi *et al.*, 2019), Katingan Cattle (Utomo, 2010), Pesisir Cattle
18 (Adrial, 2010), Indonesia local sheep (DEG Madura and Rote) (Gunawan and Sumantri,
19 2007), Ethiopia local sheep (Wagari *et al.*, 2020), Nigeria local sheep (Dauda *et al.*,
20 2018), Indonesian local buffalo (Anggraeni *et al.*, 2011; Rusdin *et al.*, 2018; Johari *et*
21 *al.*, 2009). Based on those research, it is very important to conduct a study on
22 morphological characteristic at kuantan cattle in different population There has not been
23 any report related to this matter before. The present study aimed to describe the
24 morphological characteristic and estimated genetic distance between the population of
25 Kuantan cattle from three subpopulations (Cerenti, Inuman, and Kuantan Hilir).

1 **Materials and Methods**

2 This study was conducted in Kuantan Singingi Regency. It is located at 0°00'
3 North Latitude - 1°00' South Latitude and 101°02' East Longitude - 101°55' East
4 Longitude (Figure 1).

5 **Data Collection**

6 Quantitative morphological traits were observed in 40 Kuantan male, and 150
7 Kuantan females. Location and number of the animal were determined by purposive
8 sampling. Three regions were selected, that is Cerenti, Kuantan Hilir, and Inuman. only
9 animal within the age range 2 to 3 years were considered. Five body measurement taken
10 were 1) Body Length (BL)(cm):Distance with the shoulder Joint (tuberculous humeri) to
11 the pelvic bone (tuber ischia), 2) Hip Height (HH) (cm): Vertical distance from peak hip
12 to the standing ground, 3) Wither Height (WH)(cm): wither peak, trough scapula to the
13 standing ground, 4) Chest Girth (CG) (cm): Body circumfencem behind the forelegs, 5)
14 Chest Depth (CD)(cm) (Figure 2). The age of animal was estimated by the dentition
15 method.
16

17 **Statistical Analysis**

18 Descriptive statistics (mean, SE, and CV) and t test were used to analyzed body
19 measurement. PCA was used to determine the major quantitative variable and the
20 person's coefficient correlation (r) between them and analyzed using XLSTAT (Vidal *et*
21 *al.*, 2020). The Dendogram was created based on Hierarchical Clustering Analysis (HCA)
22 using XLSTAT program.
23
24
25

Results and Discussions

Descriptive and Principle Component Analysis (PCA) of Kuantan cattle

The population of Kuantan cattle found in two regions, Kuantan Singingi and Indragiri Hulu. Cerenti, Kuantan Hilir, and Inuman located in Kuantan Singingi. This region is adjoining West Sumatera Province. Generally, Kuantan cattle have a small size like pesisir cattle in West Sumatra. These cattle were reared mainly on an extensive rearing system (Figure 3). The coat color of Kuantan cattle varied in different colors i.e. white, brown, grey, and black.

Descriptive statistics (Mean and Covariant Variance (CV) of Kuantan cattle were shown in Table 1. Chest Girth trait of Kuantan Cattle in Kuantan Hilir subpopulation, differ significantly higher than Cerenti and Inuman. Genetic and Environment has been the main factor in influenced of body measurement. This result corresponds well with Pundir *et al.*, (2015) who also find a difference significantly in body measurement between the population in three Indigenous cattle in India. Sex factors also affected body measurement. Jakaria *et al.*, (2019) reported live body weight and body measurement of Bali cattle for males are higher than females. Said *et al.*, (2017) also reported morphometric traits on Pasundan cattle males were higher than females. The minimum variability of Kuantan cattle male and female trait in wither height, and maximum variability in chest depth.

Variation of body measurement could indicate screening adaptive genetic diversity. The body measurement of kuantan cattle was observed in this study lower than some adult Indonesian indigenous and local cattle such as Bali, pasundan, pesisir, Katingan, and Madura cattle (Said *et al.*, 2017; Adrial., 2010). All variable also lower than Indigenous cattle in other countries, such as kuri cattle from Nigeria (Grema *et al.*, 2017), Begait cattle from Ethiopia (Makonnen *et al.*, 2020), Sudan Zebu Cattle from Sudan

1 (Alsiddig *et al.*, 2020), and indigenous cattle of assam from India (Keyastha *et*
2 *al.*, 2020).

3 Pearson's correlation coefficient between all variables according to Principal
4 Component Analysis (PCA) were shown in Table 2. Some values are found to be higher
5 than 0.3 and indicated a significant correlation. A significant correlation between, BL-
6 WH, WH-CG, HH-CG, and CG-CD. A highly positive correlation was found between CG-
7 CD.

8 This correlation corresponds well with Putra *et al.*, (2020) who also find a positive
9 correlation between CG and CD in Pasundan Cow (0.56). this estimate was higher
10 compare to Adinata *et al.*, (2016), who reported CG and CD correlation in Jabres cattle
11 0.247. Tyasi *et al.*, (2020) also determined the correlation between body measurement
12 and the bodyweight of Nguni cattle, and conclude there is a highly positive correlation.
13 However, score correlation between body measurement influenced by breed, age, and
14 type of animal.

15 Eigenvalues and percent of total variance along with factor loading body
16 measurement of Kuantan cattle are shown in Table 3. Three factors with eigenvalues
17 superior to 1 were observed. The first factor explained 32.77% variance effect by body
18 measurement (positively high for CG, WH, and CD). The second factor described body
19 shape contributes 25.83% of the total variability. Another study on local cattle in
20 Indonesia, such as Sulasmi *et al.*, (2017), showed body length and chest circumference
21 as an identifier body size and shape of Pasundan Bull, but in pasundan cows, the
22 identifier of body size and shapes of pasundan cows were body length and high at
23 withers. The proportion of total variance explained by the first component in this study
24 lower than Pasundan Cattle Sulasmi *et al.*, (2017); Putra *et al.*, (2020), Aceh cattle, PO,

1 and Bali (Mahmudi *et al.*, 2019), Taro white Cattle (Heryani *et al.*, 2018), but higher than
2 assam hill cattle (21.93%),

3 Animal's plot in bi-dimensional representation (F1-F2) according to PCA shown
4 in Figure 4. The bi-dimensional presentation of individuals showed that the Kuantan
5 cattle were not separated based on three populations.

6

7 **Identification of Kuantan Cattle Group using Hierarchical Cluster Analysis (HCA)**

8 The dendrogram showed that there are three clusters of Kuantan cattle based on
9 HCA in six (6) quantitative morphological trait (Figure 5). First cluster consists of 66
10 animals, mostly from Inuman sub population. Second cluster include of 79 animals,
11 mostly from Kuantan Hilir sub population, and third cluster, include of 46 animals, mostly
12 from Cerenti. Based on dendrogram, Kuantan cattle in Kuantan hilir sub population
13 have a significant distanced with Cerenti sub population.

14 Bacchouce *et al.*, (2015) also found three cluster based on HCA analysis of
15 native bovine population in Northern Tunisia, and each cluster consist of animal from
16 different populations. Eduoard *et al.*, (2019) also classified west african dwarf ewes to
17 three clusters based on eight major quantitative morphological variable. Each cluster
18 were unspecific with the three agro ecological zones of the study.

19

20 **Conclusion**

21 All the traits in Body Measurement were higher in Kuantan Hilir than that of
22 Cerenti and Inuman. PCA analysis showed the first factor explained 32.71% variance
23 effect by body measurement (Positively high for CG, WH, and CD). The second factor
24 described body shape contributes 25.87% of the total variability. The dendrogram
25 showed there are three clusters of Kuantan Cattle based on HCA Analysis.

Commented [A2]: Hasil dendrogram ini mohon dikaji/dimaknai lebih dalam. Perlu di jelaskan secara ilmiah alasan atau dugaan bahwa mengapa bisa terjadi jarak genetik yang cukup jauh antara sapi Kuantan di sub populasi Kuantan Hilir dengan di sub populasi Cerenti ? Apakah karena akibat pengaruh geografis, atau adanya penyesuaian gen asing pada salah satu sub populasi, atau pengaruh lainnya (misal adanya kebijakan pemerintah). Perlu juga dijelaskan upaya konservasi apa yang perlu dilakukan agar ke depan populasi sapi Kuantan di ketiga sub populasi tersebut tidak jauh jarak genetiknya ? Atau bagaimana upaya meningkatkan produktivitas sapi Kuantan berdasarkan informasi perbedaan jarak genetik tersebut ? Intinya, perlu kajian lebih dalam dalam memaknai hasil dendrogram.

References

- 1
2 Adinata, Y., P.D. Aryogi, M. Luthfi, A. Rasyid, and N.H. Krishna. 2016. Phenotypic
3 Characterization of Jawa Brebes (Jabres) Cattle. In Proceedings of International
4 Seminar on Livestock Production and Veterinary Technology (pp. 24-29).
- 5 Adrial . 2010. Potensi Sapi Pesisir dan Upaya Pengembangannya di Sumatera Barat.
6 Jurnal Penelitian dan Pengembangan Pertanian. 29 (2):66-72
- 7 Anggraeni, A., C. Sumantri, L. Praharani, Dudi and E. Andreas. 2011. Estimasi Jarak
8 Genetik Kerbau Rawa Lokal Melalui Pendekatan Analisis Morfologi. Indonesian J
9 Anim Vet Sci. 16(3): 199-210
- 10 Alsiddig, M.A., S.A. Babiker, M.Y. Galal, and M Ali. 2010. Phenotypic Characterization
11 of sudan zebu cattle. Reseaerch J of Anim and Vet Sci. 5 :10-17
- 12 Baccouche, R., B. Jemmal., M. Haddad., NM Hamdi, MB Hamouda. 2015. Phenotypic
13 Characterization of Native Bovine Population in Northern Tunisia. App. Sci. Report.
14 10(1):38-44
- 15 Dauda, A., H. Abbaya, and V.N. Ebegbulem. 2018. Application of Multifactorial
16 Discriminant Analysis of Morphostructural Differentiation of Sheep. Journal of
17 Genet and Genet Eng. 2(2):11-16
- 18 Edouard, N. K., B.K Lacine, K.N. Cyrille, L.N. Etienne, D.G. Kossigan, S. Mamadou, and
19 Y.G.C. Valentine. 2018. Multivariate Analysis for Morphological Characteristics of
20 N'Dama Cattle Breed in Two Agro-ecological Zones of Côte d'Ivoire. European Sci
21 J. 14(3):602
- 22 Edouard NK, KG Severin, KN Cyrille, LN Etienne, EJ Yves, DGK Charles, S Mamadou
23 and YGC Valentine. 2019. Primary Morphological Characterization of West African
24 Dwarf (Djallonké) Ewes from Côte d'Ivoire Based on Qualitative and Quantitative
25 Traits. Int. J. Genet. Mol. Biol. 11(2): 16-28

- 1 FAO. 2013. Animal Production and Health Guidelines. Molecular Genetic
2 Characterization of Animal Genetic Resources. ISAG-FAO. Rome. P.67-85
- 3 FAO. 2015. The Second Report on The State of The world's "Animal genetic resources
4 for food and agriculture. FAO Commission on Genetic Resources for Food and
5 agriculture. Rome
- 6 Grema, M., A.Traore, M. Issa, M. Hamani, M. Abdou, and I. Fernandes. 2017.
7 Morphological Assesment of Niger Kuri Cattle Using Multivariate Methods. South
8 African J of Anim Sci. 47 (4): 505-515
- 9 Gunawan and C. Sumantri. 2007. Karakteristik Morfometrik Ukuran Tubuh dan Bentuk
10 Domba Ekor Gemuk Pulau Madura dan Rote dengan Menggunakan Analisis
11 Komponen Utama. Buletin Peternakan 31(4):186-199
- 12 Heryani, L.G.S.S., N.N.W. Susari, and I.W.N.F. Gunawan. 2018. Variabel Komponen
13 Utama pada Morfometrik Sapi Putih Taro Berdasarkan Pengukuran Ukuran
14 Badan. Buletin Veteriner Udayana. 10(1):93-99
- 15 Hikmawaty, H., A. Gunawan, R.R Noor, and J. Jakaria. 2017. Identifikasi Ukuran Tubuh
16 dan Bentuk Tubuh Sapi Bali di Beberapa Pusat Pembibitan Melalui Pendekatan
17 Analisis Komponen Utama. JIPTHP 2(1):231-237
- 18 Jakaria., Sutikno, M.F. Ulum, R. Priyanto. 2019. Live body weight assessment based on
19 body measurement in bali cattle (*Bos Javanicus*) at extensive rearing system.
20 Pakistan J.Life.Soc.Sci.17(1):17-23
- 21 Johari S., E. Kurnianto, Sutopo and W.A. Hamayanti. 2009. Multivariate analysis on
22 phenotypic traits of body measurement in swamp buffalo (*bubalus bubalis*). J
23 Indonesian.Trop.Anim.Agric. 3(4):289-294

- 1 Kayastha, R.B., G. Zaman, R.N. Goswami and A. Haque. 2011. Physical and
2 Morphometric Characterization of Indigenous Cattle of Assam. *Open Vet J.* 1(1):7-
3 9
- 4 Mahmudi, M., R. Priyanto, and J Jakaria. 2019. Karakteristik Morfometrik Sapi Aceh,
5 Sapi PO dan Sapi Bali Berdasarkan Analisis Komponen Utama (AKU). *JIPTHP*,
6 7(1): 35-40.
- 7 Putra, W.P.B., F. Hilmawan, and J. Arifin. 2020. Characterization in two Indonesian Bos
8 Indocus Cattle Breeds based on Morphometric Measurements. *TJVR.* 4(1):29-32
- 9 Putra, W.B.P. 2020. The Assesment of body weight of sumba ongole cattle (*bos indicus*)
10 by body measurement. *MJAVL.* 10(1):52-57
- 11 Putra, W.P.B., S. Said, and J. Arifin. 2020. Principal Component Analysis (PCA) of Body
12 Measurements and Body Indices In The Pasundan Cow. *Black Sea J of Agr* 3(1):
13 49-55
- 14 Pundir, R.K., P.K. Singh, and D.K. Sadana. 2015. Multivariate Analysis of Morphometric
15 Traits of Three Different Indigenous Cattle Populations from North East States of
16 India. *Indonesian J Anim Vet Sci.* 20(2):79-86
- 17 Mekonnen, T and S. Meseret. 2020. Characterization of Begait Cattle using
18 morphometric and qualitative traits in western zone of Tigray Ethiopia. *Int. J.*
19 *Livest.Prod.* 11(1):21-23
- 20 Rusdin, M., D.D. Solihin, A. Gunawan, C. Talib, and C. Sumantri. 2018. Sifat Sifat
21 Kuantitatif dan Jarak Genetik Kerbau lokal Sulawesi Tenggara Berdasarkan
22 Pendekatan Morfologi. *JIPi.* 23(3):203-210
- 23 Said, S., W.P.B. Putra, S. Anwar, P.P. Agung, and H. Yuhani. 2017. Phenotipic,
24 Morphometric Characterization and population Structure of Pasundan Cattle at
25 West Java Indonesia. *Biodiversitas.* 18(4):1638-1645

- 1 Sulasmi., A. Gunawan, R. Priyanto, C. Sumantri, J. Arifin. 2017. Keseragaman dan
2 Kedekatan Morfometrik Ukuran Tubuh Sapi Pasundan. *J Veteriner*. 18(2): 263-273
- 3 Tyasi, T.L., Nhlakanipho, D. Mathye, L. Trudy, Rashijane, K. Mokoena, K.M.
4 Makgowo, M.C. Mathapo, K.M. Molabe, P.M. Bopape, and D. Maluleke. 2020.
5 Correlation And Path Analysis of Body Weight And Biometric Traits Of Nguni Cattle
6 breed. *J Adv Vet Anim Res*. 7(1): 148–155.
- 7 Utomo, BN., R.R. Noor, C. Sumantri, I. Supriatna, E.D. Gunardi. 2010. Morphometric
8 performances of Katingan cattle in Central Kalimantan. *Indonesian J Anim Vet Sci*.
9 15(3):220-230
- 10 Wagari, G., T.Getachew, E. Bayou.2020. Multivariate analysis of phenotypic traits of
11 indigenous sheep revealed new population in western part of Ethiopia. *Int J Agric*
12 *Sc Food Technol* 6(1): 050-057
- 13 Vidal, N.P., C.F. Manful, T.H. Pham, P. Stewart, P.D. Keough, and R.H. Thomas. 2020.
14 The Use of XLSTAT In Conducting Principal Component Analysis (PCA) When
15 Evaluating The Relationships Between Sensory And Quality Attributes In Grilled
16 Foods. *MethodsX*. 302(7):1-9




Figure 1 Map of Study Region





Figure 2 Scheme of Five Body Measurement in Kuantan Cattle (1) Body Length, 2)Withers Height, 3)Hip Height 4)Chest Girth, 5) Chest Depth

1
2
3
4
5




6 Figure 3 Extensive Ranging System of Kuantan Cattle



7
8
9
10

Figure 4 Bi Dimensional Representation of Individual in Three Sub Population of Kuantan Cattle (C= Cerenti, KH= Kuantan Hilir, I= Inuman)



11
12
13
14
15
16
17

Figure 6 Dendrogram of Kuantan Cattle Based on Hierarchical Classification (C: Cerenti, I: Inuman, K;Kuantan Hilir)

Commented [A3]: KH: Kuantan Hilir

1 Table 1. Descriptive Body Measurement of Kuantan Cattle

Sex	Variable	Kuantan Cattle Sub Population		
		Cerenti	Kuantan Hilir	Inuman
Male		Mean ± SE(CV) N=9	Mean ± SE(CV) N=18	Mean ± SE(CV) N=13
	Body Length	103.22 ± 2.91 (2.82)	103.78 ± 1.83 (1.77)	103.92 ± 1.75(1.69)
	Wither Height	97.44 ± 2.74 (2.82)	99.27 ± 2.74 (1.24)	98.46 ± 2.29(2.33)
	Hip Height	103.56 ± 2.88 (2.78)	103.89 ± 1.60 (1.54)	104.54 ± 3.18(3.04)
	Chest Girth	115.00 ± 8.63 ^a (7.51)	126.22 ± 4.80 ^b (3.80)	123.62± 5.80 ^b (4.16)
	Chest depth	60.22 ± 6.02 (9.99)	60.944 ± 2.27(4.40)	61.769±0.67(3.92)
Female		Mean ± SE(CV) N=40	Mean ± SE(CV) N=64	Mean ± SE(CV) N=46
	Body Length	102.28±0.30(1.89)	103.34±0.22(1.73)	103.32±0.27(1.79)
	Wither Height	97.92±0.31(2.03)	99.18±0.16(1.35)	99.63±0.23(1.59)
	Hip Height	105.13±0.33 ^a (2.01)	103.19±0.20 ^b (1.57)	103.19±0.21 ^b (1.42)
	Chest Girth	120.50±0.44 ^a (2.34)	126.14±0.53 ^a (3.37)	124.83±0.64 ^a (3.54)
	Chest depth	60.62±0.66(6.92)	62.46 ±0.30(3.88)	61.46±0.35(3.91)

N= individual number, Superscript in the same row with different superscripts are significantly different (P<0.05)

2
3 Table 2 Pearson's Correlation Overall Population Among Body Measurement of Kuantan
4 Cattle

Variables	BL	WH	HH	CG	CD
BL	1				
WH	0,300**	1			
HH	-0,063	-0,108	1		
CG	-0,052	0,371**	0,185*	1	
CD	-0,046	0,111	0,022	0,413**	1

6 BL:Body Length, WH : Wither Height, HH:Hip Height, CG: Chest Girth, CD: Chest
7 Depth


8
9 Table 3. Eigenvelues and Percent of Total Varicance Along With Factor Loading
10 Body Measurement of Kuantan Cattle

Variable	F1	F2	F3
BL	0,187	0,750	0,392
WH	0,670	0,526	0,061
HH	0,135	-0,539	0,813
CG	0,840	-0,253	0,023
CD	0,654	-0,317	-0,371
Eigenvalue	1.635	1.294	0.956
Explained Variance (%)	32.705	25.870	19.125
Cumulative Variance (%)	32.705	58.575	77.700

11 Bl: Body Length, WH : Wither Height, HH: Hip Height, CG: Chest Girth, CD: Chest
12 Depth

13

1 **Determination of Morphological Characteristics in Kuantan Cattle Using**
2 **Multivariate Analysis**
3
4



ABSTRACT

The objective of this research was to characterized morphology and estimated genetic distance between ~~intra the~~ population of Kuantan cattle. A Total of 213 cattle (44 male and 169 female with age ranging from 2-3 years) were used in this study and collected from extensive ranging systems in ~~Three~~ sub-population (Cerenti, Inuman, and Kuantan Hilir regions) Kuantan Singingi Regency, Riau Province. Five variables were measured that is Body Length (BL)_(cm), Wither Height (WH)_(cm), Hip Height (HH)_(cm), Chest Girth (CG)_(cm), and Chest Depth (CD)_(cm). Data obtained were ~~analyzed,~~ descriptive analysis, Principal Components Analysis (PCA) and Hierarchichal Clustering Analysis (HCA) using XLSTAT program. All variables of body measurement in the Kuantan Hilir region were higher than Cerenti dan Inuman, Kuantan Singingi Regency, ~~Riau Province~~. The first factor in PCA described body measurement contributed 32.77%, and the second factor described body shape contribute 25.83% of total variability. The dendrogram showed there is ~~Three~~ clusters of Kuantan Cattle based on this research.

Keywords: Kuantan cattle, PCA, Genetic Distance, Morphological Characteristics

Introduction

Indonesia has a large variety of indigenous animal genetic resources (AnGR). One of them is Kuantan cattle. Kuantan cattle are one of the local breeds in Riau Province (Indonesia) that play an important role in the maintenance of rural population living, social, religious, and traditional celebration. Kuantan cattle ~~in Riau Province~~ very adaptive in an extensive management system, and lack quality feeder. Based on data in the Kuantan Singingi government, the population of Kuantan cattle ~~decreasing from~~ 2016/2017.

Commented [A1]: Any reason about that?? Presently, How many population of Kuantan cattle? Put in this part

1 99The decrease in grazing areas and lack of bulls make the population
2 decrease. The absence of a structured and targeted breeding program in Kuantan
3 cattle, and the crossing of Kuantan cattle with other breeds without evaluation, is
4 feared genetic drift. FAO (201P5) reported states that cattle are categorized into
5 endangered groups if the total number of males is less than or equal to 20 and greater
6 than 5.

7 Conservation and production program require strategy in management and
8 detail information about animal genetic resources (FAO, 2013). Morphological
9 characterization of the breed, very important to develop the breeding scheme.
10 Improvement in the breeding system is very important to increase the population and to
11 improve the quality of Kuantan cattle.

12 The multivariate technique is one of the methods to characterize local breed
13 and has been found very suitable to identified genetic variation within and between
14 populations. (PCA) can identify the major quantitative variable, and
15 discriminant analyses more suitable in assessing to evaluate their differentiation.
16 Recently, this technique has been suitable to characterize Indonesia local breed cattle
17 (Bali cattle (Hikmawaty *et al.* 2017), Pasundan Cattle (Sulasmi *et al.*, 2017)), Ongol
18 and Sumba Ongol (Putra *et al.*, 2020) Aceh Cattle (Mahmudi *et al.*, 2019), Katingan
19 Cattle (Utomo, 2010), Pesisir Cattle (Adrial, 2010), Indonesia local sheep (DEG
20 Madura and Rote) (Gunawan and Sumantri, 2007), Ethiopia local sheep (Wagari *et al.*,
21 2020), Nigeria local sheep (Dauda *et al.*, 2018), Indonesian local buffalo (Anggraeni *et*
22 *al.*, 2011; Rusdin *et al.*, 2018; Johari *et al.*, 2009). Based on those research, it is very
23 important to conduct a study on morphological characteristic at kuantan cattle in
24 different population. There has not been any report related to this matter before. The
25 present study aimed to describe the morphological characteristic and estimated genetic

1 distance between the population of Kuantan cattle from ~~T~~three sub_populations
2 (Cerenti, Inuman, and Kuantan Hilir).

3 **Materials and Methods**

4 This study was conducted in Kuantan Singingi Regency. It is located at 0°00'
5 North Latitude - 1°00' South Latitude and 101°02' East Longitude - 101°55' East
6 Longitude (Figure 1).

7 8 **Data Collection**

9 Quantitative morphological traits were observed in 40 Kuantan male, and 150
10 Kuantan females. Location and number of the animal were determined by purposive
11 sampling. Three regions were selected, that is Cerenti, Kuantan Hilir, and Inuman. only
12 animal within the age range 2 to 3 years were considered. Five body measurement
13 taken were 1) Body Length (BL)(cm):Distance with the shoulder Joint (tuberculous
14 humeri) to the pelvic bone (tuber ischia), 2) Hip Height (HH) (cm): Vertical distance
15 from peak hip to the standing ground, 3) Wither Height (WH)(cm): wither peak, trough
16 scapula to the standing ground, 4) Chest Girth (CG) (cm): Body circumfencem behind
17 the forelegs, 5) Chest Depth (CD)(cm) (Figure 2). The age of animal was estimated by
18 the dentition method.

19 20 **Statistical Analysis**

21 Descriptive statistics (mean, SE, and CV) and t test were used to analyzed
22 body measurement. PCA was used to determine the major quantitative variable and
23 the person's coefficient correlation (r) between them and analyzed using XLSTAT
24 (Vidal *et al.*, 2020). The Dendogram was created based on Hierarchical Clustering
25 Analysis (HCA) using XLSTAT program.

Results and Discussions

Formatted: Indonesian

Descriptive and Principle Component Analysis (PCA) of Kuantan cattle

The population of Kuantan cattle found in two regions, Kuantan Singingi and Indragiri Hulu. Cerenti, Kuantan Hilir, and Inuman located in Kuantan Singingi. This region is adjoining West Sumatera Province. Generally, Kuantan cattle have a small size like pesisir cattle in West Sumatra. These cattle were reared mainly on an extensive rearing system (Figure 3). The coat color of Kuantan cattle varied in different colors i.e. white, brown, grey, and black.

Commented [A2]: What do you mean????you mean is curently??

Descriptive statistics (Mean and Covariant Variance (CV) of Kuantan cattle were shown in Table 1. Chest Girth trait of Kuantan Cattle in Kuantan Hilir subpopulation, differ significantly higher than Cerenti and Inuman. Genetic and Environment has been the main factor in influenced of body measurement. This result corresponds well with Pundir *et al.*, (2015) who also find a difference significantly in body measurement between the population in three Indigenous cattle in India. Sex factors also affected body measurement. Jakaria *et al.*, (2019) reported that live body weight and body measurement of Bali cattle for males are higher than females. Said *et al.*, (2017) also reported that morphometric traits on Pasundan cattle males were higher than females. The minimum variability of Kuantan cattle male and female trait in wither height, and maximum variability in chest depth.

Variation of body measurement could indicate screening adaptive genetic diversity. The body measurement of kuantan cattle was observed in this study lower than some adult Indonesian indigenous and local cattle such as Bali, pasundan, pesisir, Katingan, and Madura cattle (Said *et al.*, 2017; Adrial., 2010). All variable also

1 lower than Indigenous cattle in other countries, such as kuri cattle from Nigeria
2 (Grema *et al.*, 2017), Begait cattle from Ethiopia (Makonnen *et al.*, 2020), Sudan Zebu
3 Cattle from Sudan (Alsiddig *et al.*, 2020), and indigenous cattle of assam from India
4 (Keyastha *et al.*, 2020).

5 Pearson's correlation coefficient between all variables according to Principal
6 Component Analysis (PCA) were shown in Table 2. Some values are found to be
7 higher than 0.3 and indicated a significant correlation. A significant correlation
8 between, BL-WH, WH-CG, HH-CG, and CG-CD. A highly positive correlation was
9 found between CG-CD.

10 This correlation corresponds well with Putra *et al.*, (2020) who also find a
11 positive correlation between CG and CD in Pasundan Cow (0.56). this estimate was
12 higher compare to Adinata *et al.*, (2016), who reported CG and CD correlation in
13 Jabres cattle 0.247. Tyasi *et al.*, (2020) also determined the correlation between body
14 measurement and the bodyweight of Nguni cattle, and conclude there is a highly
15 positive correlation. However, score correlation between body measurement influenced
16 by breed, age, and type of animal.

17 Eigenvalues and percent of total variance along with factor loading body
18 measurement of Kuantan cattle are shown in Table 3. Three factors with eigenvalues
19 superior to 1 were observed. The first factor explained 32.77% variance effect by body
20 measurement (positively high for CG, WH, and CD). The second factor described body
21 shape contributes 25.83% of the total variability. Another study on local cattle in
22 Indonesia, such as Sulasmi *et al.*, (2017), showed body length and chest
23 circumference as an identifier body size and shape of Pasundan Bull, but in pasundan
24 cows, the identifier of body size and shapes of pasundan cows were body length and
25 high at withers. The proportion of total variance explained by the first component in this

1 study lower than Pasundan Cattle Sulasmi *et al.*, (2017); Putra *et al.*, (2020), Aceh
2 cattle, PO, and Bali (Mahmudi *et al.*, 2019), Taro white Cattle (Heryani *et al.*, 2018), but
3 higher than assam hill cattle (21.93%),

4 Animal's plot in bi-dimensional representation (F1-F2) according to PCA shown
5 in Figure 4. The bi-dimensional presentation of individuals showed that the Kuantan
6 cattle were not separated based on three populations.

7

8 **Identification of Kuantan Cattle Group using Hierarchical Cluster Analysis (HCA)**

9 The dendrogram showed that there are three clusters of Kuantan cattle based
10 on HCA in six (6) quantitative morphological trait (Figure 5). First cluster consists of 66
11 animals, mostly from Inuman sub population. Second cluster include of 79 animals,
12 mostly from Kuantan Hilir sub population, and third cluster, include of 46 animals,
13 mostly from Cerenti. Based on dendogram, Kuantan cattle in Kuantan hilir sub
14 population have a significant distanced with Cerenti sub population.

15 Bacchouce *et al.*, (2015) also found three cluster based on HCA analysis of
16 native bovine population in Northen Tunisia, and each cluster consist of animal from
17 different populations. Eduoard *et al.*, (2019) also classified west african dwarf ewes to
18 three clusters based on eight major quantitative morphological variable. Each cluster
19 were unspecific with the three agro ecological zones of the study.

20

21

21 **Conclusion**

22 All the traits in Body Measurement were higher in Kuantan Hilir than that of
23 Cerenti and Inuman.(PCA) analysis showed the first factor explained 32.71%
24 variance effect by body measurement (Positively high for CG, WH, and CD). The
25 second factor described body shape contributes 25.87% of the total variability. The

1 dendrogram showed there are three clusters of Kuantan Cattle based on HCA
2 Analysis.

3 **References**

4 Adinata, Y., P.D. Aryogi, M. Luthfi, A. Rasyid, and N.H. Krishna. 2016. Phenotypic
5 Characterization of Jawa Brebes (Jabres) Cattle. In Proceedings of International
6 Seminar on Livestock Production and Veterinary Technology (pp. 24-29).

7 **Adrial . 2010.** Potensi Sapi Pesisir dan Upaya Pengembangannya di Sumatera Barat.
8 Jurnal Penelitian dan Pengembangan Pertanian. 29 (2):66-72. (In Indonesian).

Commented [A3]: In english

Formatted: Indonesian

9 **Anggraeni, A., C. Sumantri, L. Praharani, Dudi and E. Andreas.** 2011. Estimasi Jarak
10 Genetik Kerbau Rawa Lokal Melalui Pendekatan Analisis Morfologi. Indonesian J
11 Anim Vet Sci. 16(3): 199-210. (In Indonesian).

Commented [A4]: In english

Formatted: Indonesian

12 Alsiddig, M.A., S.A. Babiker, M.Y. Galal, and M Ali. 2010. Phenotypic Characterization
13 of sudan zebu cattle. Reseaerch J of Anim and Vet Sci. 5 :10-17

14 Baccouche, R., B. Jemmal., M. Haddad., NM Hamdi, MB Hamouda. 2015. Phenotypic
15 Characterization of Native Bovine Population in Northern Tunisia. App. Sci.
16 Report. 10(1):38-44

17 Dauda, A., H. Abbaya, and V.N. Ebegbulem. 2018. Application of Multifactorial
18 Discriminant Analysis of Morphostructural Differentiation of Sheep. Journal of
19 Genet and Genet Eng. 2(2):11-16

20 Edouard, N. K., B.K Lacine, K.N. Cyrille, L.N. Etienne, D.G. Kossigan, S. Mamadou,
21 and Y.G.C. Valentine. 2018. Multivariate Analysis for Morphological
22 Characteristics of N'Dama Cattle Breed in Two Agro-ecological Zones of Côte
23 d'Ivoire. European Sci J. 14(3):602

24 Edouard NK, KG Severin, KN Cyrille, LN Etienne, EJ Yves, DGK Charles, S Mamadou
25 and YGC Valentine. 2019. Primary Morphological Characterization of West

1 African Dwarf (Djallonké) Ewes from Côte d'Ivoire Based on Qualitative and
2 Quantitative Traits. Int. J. Genet. Mol. Biol. 11(2): 16-28

3 FAO. 2013. Animal Production and Health Guidelines. Molecular Genetic
4 Characterization of Animal Genetic Resources. ISAG-FAO. Rome. P.67-85

5 FAO. 2015. The Second Report on The State of The world's "Animal genetic resources
6 for food and agriculture. FAO Commission on Genetic Resources for Food and
7 agriculture. Rome

8 Grema, M., A.Traore, M. Issa, M. Hamani, M. Abdou, and I. Fernandes. 2017.
9 Morphological Assesment of Niger Kuri Cattle Using Multivariate Methods. South
10 African J of Anim Sci. 47 (4): 505-515

11 Gunawan and C. Sumantri. 2007. Karakteristik Morfometrik Ukuran Tubuh dan Bentuk
12 Domba Ekor Gemuk Pulau Madura dan Rote dengan Menggunakan Analisis
13 Komponen Utama. Buletin Peternakan 31(4):186-199. *(In Indonesian)*,

14 Heryani, L.G.S.S., N.N.W. Susari, and I.W.N.F. Gunawan. 2018. Variabel Komponen
15 Utama pada Morfometrik Sapi Putih Taro Berdasarkan Pengukuran Ukuran
16 Badan. Buletin Veteriner Udayana. 10(1):93-99. *(In Indonesian)*,

17 Hikmawaty, H., A. Gunawan, R.R Noor, and J. Jakaria. 2017. Identifikasi Ukuran
18 Tubuh dan Bentuk Tubuh Sapi Bali di Beberapa Pusat Pembibitan Melalui
19 Pendekatan Analisis Komponen Utama. JIPTHP 2(1):231-237. *(In Indonesian)*,

20 Jakaria., Sutikno, M.F. Ulum, R. Priyanto. 2019. Live body weight assessment based
21 on body measurement in bali cattle (Bos Javanicus) at extensive rearing system.
22 Pakistan J.Life.Soc.Sci.17(1):17-23

23 Johari S., E. Kurnianto, Sutopo and W.A. Hamayanti. 2009. Multivariate analysis on
24 phenotypic traits of body measurement in swamp buffalo (*bubalus bubalis*). J
25 Indonesian.Trop.Anim.Agric. 3(4):289-294

Commented [A5]: In english

Formatted: Indonesian

Commented [A6]: In english

Formatted: Indonesian

Commented [A7]: In english

Formatted: Indonesian

Formatted: Font: Italic

Formatted: Font: Italic

- 1 Kayastha, R.B., G. Zaman, R.N. Goswami and A. Haque. 2011. Physical and
2 Morphometric Characterization of Indigenous Cattle of Assam. Open Vet J.
3 1(1):7-9
- 4 Mahmudi, M., R. Priyanto, and J Jakaria. 2019. Karakteristik Morfometrik Sapi Aceh,
5 Sapi PO dan Sapi Bali Berdasarkan Analisis Komponen Utama (AKU). JIPTHP,
6 7(1): 35-40. (In Indonesian)
- 7 Putra, W.P.B., F. Hilmawan, and J. Arifin. 2020. Characterization in two Indonesian
8 Bos Indocus Cattle Breeds based on Morphometric Measurements. TJVR.
9 4(1):29-32
- 10 Putra, W.B.P. 2020. The Assesment of body weight of sumba ongole cattle (bos
11 indicus) by body measurement. MJAVL. 10(1):52-57
- 12 Putra, W.P.B., S. Said, and J. Arifin. 2020. Principal Component Analysis (PCA) of
13 Body Measurements and Body Indices In The Pasundan Cow. Black Sea J of
14 Agr 3(1): 49-55
- 15 Pundir, R.K., P.K. Singh, and D.K. Sadana. 2015. Multivariate Analysis of
16 Morphometric Traits of Three Different Indigenous Cattle Populations from North
17 East States of India. Indonesian J Anim Vet Sci. 20(2):79-86
- 18 Mekonnen, T and S. Meseret. 2020. Characterization of Begait Cattle using
19 morphometric and qualitative traits in western zone of Tigray Ethiopia. Int. J.
20 Livest.Prod. 11(1):21-23
- 21 Rusdin, M., D.D. Solihin, A. Gunawan, C. Talib, and C. Sumantri. 2018. Sifat Sifat
22 Kuantitatif dan Jarak Genetik Kerbau lokal Sulawesi Tenggara Berdasarkan
23 Pendekatan Morfologi. JIPI. 23(3):203-210. (In Indonesian)

Commented [A8]: In english

Formatted: Indonesian

Commented [A9]: In english

Formatted: Indonesian

- 1 Said, S., W.P.B. Putra, S. Anwar, P.P. Agung, and H. Yuhani. 2017. Phenotypic,
2 Morphometric Characterization and population Structure of Pasundan Cattle at
3 West Java Indonesia. *Biodiversitas*. 18(4):1638-1645
- 4 **Sulasm**., A. Gunawan, R. Priyanto, C. Sumantri, J. Arifin. 2017. Keseragaman dan
5 Kedekatan Morfometrik Ukuran Tubuh Sapi Pasundan. *J Veteriner*. 18(2): 263-
6 273. (In Indonesian)
- 7 Tyasi, T.L., Nhlakanipho, D. Mathye, L. Trudy, Rashijane, K. Mokoena, K.M.
8 Makgowo, M.C. Mathapo, K.M. Molabe, P.M. Bopape, and D. Maluleke. 2020.
9 Correlation And Path Analysis of Body Weight And Biometric Traits Of Nguni
10 Cattle breed. *J Adv Vet Anim Res*. 7(1): 148–155.
- 11 Utomo, BN., R.R. Noor, C. Sumantri, I. Supriatna, E.D. Gunardi. 2010. Morphometric
12 performances of Katingan cattle in Central Kalimantan. *Indonesian J Anim Vet*
13 *Sci*. 15(3):220-230
- 14 Wagari, G., T.Getachew, E. Bayou.2020. Multivariate analysis of phenotypic traits of
15 indigenous sheep revealed new population in western part of Ethiopia. *Int J Agric*
16 *Sc Food Technol* 6(1): 050-057
- 17 Vidal, N.P., C.F. Manful, T.H. Pham, P. Stewart, P.D. Keough, and R.H. Thomas. 2020.
18 The Use of XLSTAT In Conducting Principal Component Analysis (PCA) When
19 Evaluating The Relationships Between Sensory And Quality Attributes In Grilled
20 Foods. *MethodsX*. 302(7):1-9

Formatted: Highlight

Formatted: Indonesian




Figure 1 Map of Study Region




Figure 2 Scheme of Five Body Measurement in Kuantan Cattle (1) Body Length, 2)Withers Height, 3)Hip Height 4)Chest Girth, 5) Chest Depth

1
2
3
4
5



6 Figure 3 Extensive Ranging System of Kuantan Cattle

7
8
9
10




Figure 4 Bi Dimensional Representation of Individual in Three Sub Population of Kuantan Cattle (C= Cerenti, KH= Kuantan Hilir, I= Inuman)

11
12
13
14
15
16
17




Figure 6 Dendrogram of Kuantan Cattle Based on Hierarchical Classification (C: Cerenti, I: Inuman, K:Kuantan Hilir)

1 Table 1. Descriptive Body Measurement of Kuantan Cattle

Sex	Variable	Kuantan Cattle Sub Population		
		Cerenti	Kuantan Hilir	Inuman
Male		Mean ± SE(CV) N=9	Mean ± SE(CV) N=18	Mean ± SE(CV) N=13
	Body Length	103.22 ± 2.91 (2.82)	103.78 ± 1.83 (1.77)	103.92 ± 1.75(1.69)
	Wither Height	97.44 ± 2.74 (2.82)	99.27 ± 2.74 (1.24)	98.46 ± 2.29(2.33)
	Hip Height	103.56 ± 2.88 (2.78)	103.89 ± 1.60 (1.54)	104.54 ± 3.18(3.04)
	Chest Girth	115.00 ± 8.63 ^a (7.51)	126.22 ± 4.80 ^b (3.80)	123.62± 5.80 ^b (4.16)
	Chest depth	60.22 ± 6.02 (9.99)	60.944 ± 2.27(4.40)	61.769±0.67(3.92)
Female		Mean ± SE(CV) N=40	Mean ± SE(CV) N=64	Mean ± SE(CV) N=46
	Body Length	102.28±0.30(1.89)	103.34±0.22(1.73)	103.32±0.27(1.79)
	Wither Height	97.92±0.31(2.03)	99.18±0.16(1.35)	99.63±0.23(1.59)
	Hip Height	105.13±0.33 ^a (2.01)	103.19±0.20 ^b (1.57)	103.19±0.21 ^b (1.42)
	Chest Girth	120.50±0.44 ^a (2.34)	126.14±0.53 ^a (3.37)	124.83±0.64 ^a (3.54)
	Chest depth	60.62±0.66(6.92)	62.46 ±0.30(3.88)	61.46±0.35(3.91)

N= individual number, Superscript in the same row with different superscripts are significantly different (P<0.05)

2
3 Table 2 Pearson's Correlation Overall Population Among Body Measurement of
4 Kuantan Cattle

Variables	BL	WH	HH	CG	CD
BL	1				
WH	0,300**	1			
HH	-0,063	-0,108	1		
CG	-0,052	0,371**	0,185*	1	
CD	-0,046	0,111	0,022	0,413**	1

6 BL:Body Length, WH : Wither Height, HH:Hip Height, CG: Chest Girth, CD:
7 Chest Depth

8
9 Table 3. Eigenvelues and Percent of Total Varicance Along With Factor Loading
10 Body Measurement of Kuantan Cattle

Variable	F1	F2	F3
BL	0,187	0,750	0,392
WH	0,670	0,526	0,061
HH	0,135	-0,539	0,813
CG	0,840	-0,253	0,023
CD	0,654	-0,317	-0,371
Eigenvalue	1.635	1.294	0.956
Explained Variance (%)	32.705	25.870	19.125
Cumulative Variance (%)	32.705	58.575	77.700

11 BL: Body Length, WH : Wither Height, HH: Hip Height, CG: Chest Girth, CD:
12 Chest Depth

13